

Thesis/
Reports
Jameson,
D. A.

SOUTHWESTERN CHAPARRAL, PINYON-JUNIPER
WOODLAND, AND FOREST VEGETATION TYPE

18.1

(A Problem Analysis)

by

DONALD A. JAMESON

UNITED STATES GOVERNMENT

Memorandum

1705

TO : Director

SEP 5 1967

4210
DATE: August 14, 1967


FROM : Donald A. Jameson, Project Leader

SUBJECT: Range and Wildlife Habitat Programs (Problem Analysis)

EAR

The attached problem analysis was completed prior to the recent change in the specified procedure for submitting such analyses. Although this report is no longer required, it may prove to be of value as a working tool and will be kept on file in Flagstaff for reference.

Attachments-1



SOUTHWESTERN CHAPARRAL, PINYON-JUNIPER WOODLAND,

AND FOREST VEGETATION TYPE

(A Problem Analysis)

by

DONALD A. JAMESON

Principal Plant Physiologist

Rocky Mountain Forest and Range Experiment Station

Forestry Sciences Laboratory

Flagstaff, Arizona

THE PROJECT AREA

The Southwestern chaparral, pinyon-juniper woodland, and forest vegetation types form a major portion of the western range comprising some 55 million acres (fig. 1). Although broadly grouped into three major vegetation types, the area encompasses climate conditions from semidesert to subalpine. The seasonal distribution of precipitation varies from Mediterranean to summer monsoon. Soils vary from incompletely decomposed sandstones to very heavy clays, and plant communities range from grassland to climax forest.

Grazing management ranges from sophisticated to crude. The emphasis in range improvement practices in the past has been on agronomic practices such as reseeding and brush control. Recently more attention is being paid to rotation and deferment systems.

Application of existing knowledge in the pine and pinyon-juniper types is beginning to provide a higher level of range productivity; such improvements can be expected to increase in the future. In these areas there is often an apparent conflict between range production and other uses, particularly between pine regeneration and range forage production. In the chaparral type, there is much less background information and the research approach will necessarily be different.

Thus, the problems to be considered are:

1. In the pine and pinyon-juniper types specific management and conflict of use problems which remain following management of ranges according to the best present techniques.
2. In the chaparral type the collection of ample background information to develop a more precise research program in the future.

APPROACH TO THIS ANALYSIS

The approach used in this analysis is one of a system analysis. This analysis considers that range management is a series of related systems whereby atmospheric gases, solar energy precipitation and parent material are processed to produce income to a ranch operator.

The Range System consists of four processing units:

1. The environmental integrator (soil).
2. The stationary biological processor (higher plants, lower plants, and lower animals).
3. The movable biological processor (higher animals).
4. The economic processor.

The basic system is outlined in Figure 2. Each unit has inputs from the preceeding unit and outputs to the succeeding unit. Each unit also has losses which may be utilized by other systems, but nevertheless are losses to the range system. Each unit has feedback to preceeding units. This system is further complicated by additional factors:

1. The processing units can be replaced. For example, reseedling is a fundamental change in the stationary biological processor. A change from cow-calf to steers is a fundamental change in the movable biological processor.

THE RANGE SYSTEM




Figure 2.

2. The entire system operates through time, and outputs are generally nonlinear in time. For example, net radiation, available soil moisture, consumed forage, and animal products are all cyclic. The losses are also nonlinear in time.

3. Feedback loops are generally nonlinear, although some are quasi-linear. Overlapping feedback loops occur. In spite of these complications, following through such a system does point out some fruitful areas of research. We have an understanding of the general nature of most of the common processor units, but discrepancies in input/output ratios point out a few incongruities.

PRESENT KNOWLEDGE

I. Environmental Factors

A. Precipitation

The most prominent and unique feature of the Southwestern climate is the two distinct rainfall periods. Across Arizona and western New Mexico there is a gradual reduction of summer rainfall from east to west, and a decrease in winter rainfall from west to east. This pattern results in the maximum development of the two-season rainfall patterns in the Flagstaff-Prescott area (fig. 3).

The greatest winter rainfall in Arizona comes from storms moving eastward from lower California (fig. 4). As these storms move eastward they drop most of their moisture on the desert mountains of southwestern and central Arizona. These storms are particularly noticeable along the Mogollon Rim of central Arizona. The maximum moisture from these storms is near the Sierra Ancha Experimental Forest. Downwind from the Rim there is a distinct rainshadow, which is especially pronounced in the Little Colorado River basin. The importance of these California storms is almost entirely lost by the Arizona-New Mexico border. Thus, climatically speaking, the most prominent feature in Arizona is the Mogollon Rim, with a winter wet climate on the windward side and a winter dry climate on the lee side.




Figure 3.--Two-season effect in southwestern rainfall. Two-season effect is largest in darkest areas.




Figure 4. Distribution of winter storms in the Southwest. Isohyets show rainfall at the 5,000 foot level. Dashed line shows equal March-April precipitation--actual Great Basin climate extends to the south of dashed line.

Northern Arizona and New Mexico have a winter climate very similar to the winter climate of the Great Basin. This differs from the southern California pattern in that April is a higher rainfall month than March. Figure 4 shows the line of equal March-April precipitation. Below that line the winter moisture regime is predominantly Californian; above the line it is predominantly Great Basin in nature. Generally, the Great Basin storms lose their effectiveness before reaching the Little Colorado River.

Summer precipitation is predominantly from the Sonoran summer monsoon. In Arizona the peak activity of the summer monsoon is in the south central portion (fig. 5). The effect of the summer monsoon decreases rapidly toward the north.

Since the monsoon air masses do not show distinct frontal characteristics, there is no rain shadow effect as with the winter storms. Because much of the precipitation falls as a result of convection storms, high precipitation coincides with points of abrupt topography. The highest amounts of summer precipitation is at about the same location as the highest amounts of winter precipitation, i.e., in the Payson-Sierra Ancha area.

In east central New Mexico the Great Plains pattern of precipitation becomes evident. In the Great Plains May and June are peak precipitation months, in Arizona May and June




Figure 5. Distribution of rainfall from the summer monsoon.

Solid lines are 5,000 foot isohyets. Dashed lines show ratio of May/June precipitation.

are drought months with peak precipitation being in August. Figure 5 shows the decline of May precipitation across eastern New Mexico. In west-central New Mexico the Great Plains pattern is almost completely lost. The dominant influences for western New Mexico and most of Arizona are these convection storms from the summer monsoon and winter frontal storms moving eastward from southern California.

Precipitation of various locations is shown in table 1 corresponding figures (6-10) show representative vegetation at these locations. Under each precipitation regime the vegetation grades to various species of juniper trees with increasing elevation and precipitation, but the associated species such as semidesert shrub, chaparral, sagebrush, or grasses remain as understories. All of the understories except southern desert shrub are also represented at even higher elevations where the juniper gives way to ponderosa pine.

These regimes are important in applying management practices. Techniques which are critically controlled by climate are often successful under one climatic regime, but not under another.

Climatic summaries for the southwest can be found in U.S. Weather Bureau annual summaries, in Smith (1956), Green and Sellers (1964), and Hastings (1964). Colton (1958) reported local precipitation near the San Francisco Peaks, and Thornthwaite (1942) discussed climate of the Polacca Wash. Recent rainfall distribution maps have been prepared by the U.S. Weather Bureau. Current research is being conducted by the Institute of Atmospheric



**Figure 6. Sagebrush type with predominately winter precipitation
(Fredonia-Jacobs Lake Road).**



Figure 7. Chaparral vegetation near Prescott Airport, Arizona.



Figure 8. Semidesert vegetation near Winston, New Mexico.



Figure 9. Blue grama-juniper type near Snowflake, Arizona.



Figure 10. Blue grama type near St. Johns, Arizona.

**Table 1. Precipitation and elevation at selected stations
in Arizona and New Mexico**

| Elevation | Mount Trumbull, Ariz. | Prescott, Ariz. | Winston, New Mex. | Snowflake, Ariz. | St. Johns, Ariz. |
|--|--------------------------------------|----------------------------|------------------------------|-----------------------------|-----------------------------|
| | 5,560 | 5,410 | 6,200 | 5,642 | 5,730 |
| - - - - - Precipitation - - - - - | | | | | |
| Annual | 12.50 | 12.23 | 12.14 | 11.69 | 11.39 |
| Jan. | .96 | 1.01 | .55 | .84 | .69 |
| Feb. | 1.06 | 1.08 | .60 | .62 | .70 |
| Mar. | 1.04 | .73 | .41 | .60 | .75 |
| Apr. | .73 | .69 | .45 | .63 | .55 |
| May | .53 | .24 | .52 | .48 | .52 |
| June | .40 | .32 | .82 | .49 | .60 |
| July | 1.80 | 2.08 | 2.25 | 1.87 | 2.19 |
| Aug. | 2.04 | 2.66 | 2.75 | 2.50 | 2.11 |
| Sept. | 1.08 | 1.19 | 1.95 | 1.41 | 1.28 |
| Oct. | 1.03 | .73 | .97 | 1.02 | .91 |
| Nov. | .77 | .50 | .30 | .58 | .41 |
| Dec. | 1.06 | 1.00 | .57 | .65 | .68 |
| June - Sept. | 5.32 | 6.25 | 7.77 | 6.27 | 6.18 |
| Oct. - May | 7.18 | 5.98 | 4.35 | 5.42 | 5.21 |
| Fig. No. | 6 | 7 | 8 | 9 | 10 |

Physics of Tucson, Arizona, and the U.S. Weather Bureau at Tucson, Arizona. A report by Bryson (1957) is representative of the former and a report by Kanglieser and Green (1965) is a representative of the latter. This research has not as yet related climate to vegetation.

Vishen (1954) provides a good background of climatic patterns in the United States and Baker (1944) and Geiger (1965) discusses local modification due to topography.

The effect of weather on forage plant growth in the pinyon-juniper type has been studied briefly and reported by Jameson (1965). Nelson (1934) reported the loss of grass stands due to drought on the Jornada Experimental Range; Coupland (1958) and Albertson, Tomanek, and Riegel (1957) discussed the effects of drought on the Great Plains. These studies are most useful in pointing out that drought does have severe effects on range production. There have also been reports by Upson (1937) and Lister and Schumacher (1937).

Recently, tree ring studies (Glock 1961, Fritts 1965) have shown the relationship of present climate to long term climate. The study by Fritts is particularly inclusive. This study shows that a severe drought period occurred in Arizona and western New Mexico about 1905, followed by a wet period from 1910 to 1925. Hastings (1959) concluded that loss of forage due to overgrazing during drought periods, followed by wet periods, is the most likely reason for the great increase in woody plots which has occurred in the Southwest. Humphrey (1958) reached the conclusion that the main factor was fire, but the preponderance of the evidence seems to support Hastings' conclusion.

For the period 1931-1940, according to Fritts (1965) the Arizona and western New Mexico area had growing conditions almost identical to the 1501-1940 average. The very recent conditions have not been greatly different than the 1931-40 period. The present drought in the Southwest (Searles 1951, Barr 1956) is not unusual when the climate over hundreds of years is considered; rather "drought" in the Southwest is the usual state of affairs.

Vegetation distribution maps for the Southwest have been prepared by Nicol (1952), Castetter (1956), Whiting (1942) and U.S.D.A. Forest Service (1965). Merriam (1890) first pointed out the changes in plant communities with changes in elevation, and measurements of different weather conditions in the Southwest were first reported by Pearson (1931). One of the most recent studies of climate and vegetation is that of Johnsen (1962) who discussed the relationship of climate to one-seed juniper.

In summary, there is a considerable background of precipitation information for the Southwest and more is being developed by agencies primarily concerned with weather research. Long term weather records are scarce, but adequate interpolation can usually be obtained from existing stations when a picture of the long term climate is needed. Weather observations still need to be made in conjunction with certain types of vegetation studies. Studies of probabilities of rainfall will need to be conducted if probabilistic approaches to analysis of rainfall-dependent range improvements are undertaken (Pattison 1965, Glover and LeBaron 1966).

B. Solar Radiation and Temperature

Solar radiation and temperatures are excessive in the lower regions of project area and at least adequate in all regions. There are two noticeable examples: Muhlenbergia porteri and Poa fendleriana are commonly found growing only in the shade of protective plants. Such phenomena should be documented, but it is doubtful if a feasible means of reducing solar radiation and soil temperatures can be found.

C. Gases

In rapidly growing crop land there is often a deficiency of carbon dioxide. Because of the low growth rates in the project area there apparently is no deficiency of carbon dioxide available to the plants.

II. Geology and Soils

Because of the arid climate, parent material has a very great influence on soils of the Southwest, and geologic maps are available for both Arizona and New Mexico. In addition, soil surveys by the National Forests and by the Soil Conservation Service are progressing rapidly and many survey reports should be published within the next few years. Some published information on soils and geology can be found in Martin and Fletcher (1943), Rich (1962), McKee (1936), Robinson (1913), Wilson (1962), Veatch (1920), and Carter and Cory (1932).

Approximate averages of parent materials within the Arizona pinyon-juniper type have been tabulated as follows:

| <u>Parent material</u> | <u>Thousands of acres</u> |
|------------------------|---------------------------|
| Volcanic | 4,454 |
| Kaibab formation | 3,502 |
| Jurassic sandstone | 1,411 |
| Mesa Verde formation | 1,173 |
| Coconino sandstone | 1,156 |
| Supai sandstone | 612 |
| Redwall limestone | 578 |
| Granite | 408 |
| Sand and gravel | 510 |

In the ponderosa pine type about three-fourths of the area is basalt parent material, about one-fifth is Kaibab limestone, and the remainder is divided among the other parent materials listed for the juniper type.

In the chaparral the foremost parent materials are granite and granitic outwash, with schist and basalt next in importance. These four parent materials make up about 90 percent of the type, the chaparral in general is poorly developed on basalt and best developed on granite and schist. Other parent materials include diabase, Mescal limestone, quartzite, sandstone, etc., but none of these make up more than a small percentage of the type, except in the localized area near Roosevelt Lake.

Several basalt pine soils have been studied by Clary, Ffolliott, and Zander (1966). On these soils, highest productivity was in the swales and lowest was in the rocky ridges. The same general pattern holds in the juniper soils. In addition, skeletal limestone soils produce less than skeletal soils from other parent materials in the pinyon-juniper types. This may be related to lowered phosphorus availability in these soils (Sengupta and Cornfield 1963). The possibility of greatly improving production on these rocky soils seems extremely remote.

In the juniper type the Springerville soil constitutes a problem which possibly can be solved (fig. 11). This soil is a very heavy basaltic clay which is unstable (Vertisol). With a grass cover production is not much different than other ordinary upland soils. With the presence of juniper trees, however, the production on this soil is reduced much more than production of other soils. The reasons for this decline in production are at present uncertain, but there are several promising leads. Johnsen, Cady and James (1962) indicate that the clay of the Springerville series is predominantly montmorillonite complexed with organic matter. The bulk density is high and aeration is correspondingly low. Drainage is poor. Some counts of microbial population gave 700 colonies per cc in Springerville compared to 5,000 colonies per cc in other basalt soils. A low microorganism count is typical for



Figure 11. Springerville soil series in a Utah juniper stand.

montmorillonite clays (Stotzky, 1962). All of this information indicates that the growth inhibitors present in juniper (Jameson, 1961), which are normally oxidized by microorganisms or leached by water are accumulated in the soil. The enzymes responsible for oxidation of the inhibiting chemicals are copper containing proteins (Schubert, 1965). Preliminary analyses have shown that the copper content of Springerville soils is low.

Since the Springerville soils are heavy clays, they may have some physical effects on plants because of frost heaving (Anderson 1947, Biswell 1953, Schram 1958).

III. Losses from the soil and Trampling Damage

Of the environmental factors which enter the soil complex, only soil moisture is limiting enough that we should concern ourselves with its loss. Hydrologists utilize the technique of soil moisture budgets to study such losses; extensive hydrologic studies are underway in the project area by Watershed Research and Forest Economics Research. Further work on soil moisture by Range Management Research will very likely be needed in the future.

Erosion of the soil itself is also a loss that needs to be considered. In most range management systems protection against erosion and maximizing the forage production are nearly synonymous. In the chaparral type, however, there may be a different problem. The bulk of the forage production is from shrubs which seem to provide little in the way of protection from erosion (fig. 12). Studies by Rich (1956) and Rich and



Figure 12. Soil erosion in the chaparral. Deep gullies have formed in bottom land alluvium.

Reynolds (1963) indicate that if grazing is light enough that no change in the grass cover is induced there is correspondingly no change in water yield and erosion. We have no knowledge of the erosion that could be expected if chaparral management were based on utilization of the shrub species, but we expect that losses would be mostly due to trampling rather than to reduction of cover.

IV. Decomposing Plants

In both the pine and pinyon-juniper type decomposed tree needles are a detriment to forage production (Jameson 1961, Clary and Ffolliott 1966). Although the effect is evident (fig. 13) the precise nature of the problem is still unclear. To date our attempts to identify the active components have not been successful.

V. Descriptions of Plants and Plant Communities

A. Communities

Studies on descriptions of plant communities have been confined mostly to the pinyon-juniper type as the following list of references indicate:

General.--Cottle 1931, 1932; Daubenmire 1943; Loftfield 1923-24; Lowe 1961; Rasmussen 1941; Reynolds 1962; Shantz and Zon 1924; Shreve 1942; Society of American Foresters 1957.

Pinyon-juniper.--Billings 1951, 1954; Bostick 1947; Bray 1940; Buchanon 1961; Deaver and Haskell 1955; Driscoll 1962; Eastwood 1919; Emerson 1932; Erdman 1962; Jameson, Williams, and Wilton 1962; Halls 1955; Livingston 1947; Merkle 1952;



Figure 13. Effect of pine litter on understory vegetation.

Mead 1930; Miller 1921; Olsen and Dykshuis 1940; Palmer 1920; Parker 1945; Phillips and Mulford 1912; Plummer, Rixon, and Dodwell 1904; Potter 1957; Randles 1949; Read 1915; Reveal 1944; Stagner 1962; Turner 1913; Woodbury 1933; 1947; Woodhouse 1953; Woodin 1953; Woodin and Lindsey 1954.

Pine.--Arnold 1955; Dodge 1963.

Chaparral.--Hansen 1954; Illige 1954; Pase and Pond 1962; Pond and Cable 1962.

This list is by no means complete, but it does demonstrate the wide background of basic community description available in the pinyon-juniper type in contrast to the other vegetation type. Part of the reason is the extent of the pinyon-juniper type; there are some 60 million acres compared to 4 million acres each for the chaparral and southwestern ponderosa pine. Although the problems are less extensive in the latter two types, they are just as intensive. The Arizona chaparral, for example, is made up of many subtypes (fig. 12). The fact of the diversity is recognized, but published descriptions dealing with this diversity are almost completely lacking and there is no research underway. Our present study area at Tonto Springs is located on granitic outwash and is almost pure shrub live oak. A similar study on granite or schist parent material with a more diverse plant population needs to be established.

In spite of wide scope of information in the pinyon-juniper type, one can quickly question its value in a rigorous analysis of the problem. Descriptive ecology commonly only documents

the obvious. The real problems of the ecosystems are usually not touched in such documentation. For example, soil microorganisms, soil fertility, soil plant water relationships, etc., are usually not at all considered. Management practices cannot be intelligently planned based on a description of the ecosystem, but can be planned only by being informed of the inputs available to and outputs demanded from it.

Nevertheless, without such background information there is little direction in a research program or management planning, even though the exact use of community ecology information is often not readily seen.

VI. Individual Species Components of the Plant-Lower Animal Community

We have, or soon will have, completed reports on phenology of most of our important plants. The pinyon-juniper phenology has been published, the phenology of the pine-type grasses has been submitted for publication, and phenology of the shrubs is being prepared. We also have long term records of community changes in the pine and juniper type.

In any project area the number of autecological studies that could be undertaken on individual species are practically without limit. For example, we could study light, moisture, temperature and mineral requirements for germination and growth of any number of species. In any project area we could maintain that such information is scarce, as well it would be. The southwestern chaparral, woodland, and forest ranges are no exception. A few studies have been made, for example:

Olmstead 1952. Photoperiodism of gramas.

Benedict 1941. Growth of blue grama in reduced light.

Moore 1963. Germination inhibitors in achenes of Cerocarpus.

Lister and Schumacher 1957. Precipitation and grass growth.

There are a few other examples that could be cited, but not many more.

Since the study opportunities are essentially without limit, we must restrict ourselves in some way. First, which species are most important? Generally this means which are the most widespread, but areal extent may not be the only criteria. A species which is ordinarily produced in a fair amount and utilized in porportion by grazing animals does not demand the attention that a widespread weed species or a scarce, but highly utilized species does. Also, a limited, but preferred species loses much of its importance when a more widespread species is an adequate substitute. Secondly, what factors of importance to a species can be controlled? For example, most of our species could be vastly improved by additional water, but the possibilities of furnishing additional water are extremely remote.

Because of these factors, the choices of autecological studies are highly subjective. In our case we have chosen snakeweed (Gutierrezia spp.) (fig. 14) as the species demanding the most attention, and we have chosen it for the following reasons:

Not at all
after grass

(1) It is extremely widespread. Gutierrezia is the second most important genus in the pinyon-juniper type, exceeded only by Juniperus, and it also extends into the pine and chaparral types.

(2) It is apparently not utilized at all by cattle under normal management even though it is utilized under very heavy grazing.

(3) It exhibits extreme variations in abundance between years. This indicates that there may be some facet of its ecology which would provide a clue to its more effective management.

(4) Although it is relatively easy to control with chemicals, we have no knowledge of how to stop or even reduce reinvasions.

Because of these reasons, intensive autecological research should begin with Gutierrezia. Background research includes Campbell and Bomberger (1943), a thesis from the University of Arizona, and unpublished research by Thomas N. Johnsen, Jr., ARS. In addition to its plant ecological importance, snakeweed is suspected of causing abortion.

VII. Plant Losses to the Range System

A plant loss to the range system is a species or parts of species which are not consumable, the retention of which is not required to sustain the desired plant community. Which species or parts of species are not consumed varies with the type of range management applied, but in general we can divide species into three general groups:



Figure 14. Gutierrezia is the most widespread genus in the project area.

(1) Those of which a major portion is consumed under most normal management systems. These are the "ice cream" and "key" species.

(2) Those which are not consumed at all or at least not under any but the most severe grazing pressure. In our project area this included crucifixion thorn, snakeweed, prickly pear, and large trees.

(3) Species which the degree of utilization can be regulated by type of management applied. Such species as shrub live oak and tobosa are in this category.

Ideally we should measure the production of both the consumable and nonconsumable species in order to determine the output of the plant community system. Practically, however, this is difficult to do and may be misleading even if it is done. For example, what comparisons can be drawn between the production of one pound of juniper and one pound of grass? What comparisons can be drawn between one pound of a species which grows mostly in the spring and a pound of a species which grows mostly in the summer? Because of these and similar difficulties, it is just as appropriate and much more feasible to measure the nonconsumable plants in terms which are easy to measure, and measure only the consumable plants in such output units as pounds. We find that relationships drawn between consumable plants expressed as weight and nonconsumable plants expressed in some other units are very

helpful. Examples are those of the chaparral (Pond 1964) the pinyon-juniper (Arnold, Jameson, and Reid 1964) and the pine (Pearson 1954). In each case there is a normal relationship between species or classes of species--a departure from the normal indicates an unusual condition. The Springerville soil in the juniper type is a case in point. In these soils the herbage reduction associated with a given juniper tree cover is much greater than for other soils (fig. 15); this indicates that a different set of factors are operating.

Usually we think that trees in semiarid situations represent losses to the range system because of competition for soil moisture with the consumable species. Sometimes, however, this is not the case. For example, there is apparently no competition for soil moisture between blue grama and juniper, rather the effect seems to be chemical or physical. Other examples of noncompetitive effects for both the project area and other areas are found in the following references:

Arnold 1964; Börner 1960; Cacini 1963; Evanari 1957; Garb 1961; Glendenning 1939; Grummer 1961; Jameson 1961; Muller 1953; Muller and Muller 1956; Rovira 1962; Sumere 1960; Woods 1960.

Examples of chemical effects of one species upon another are so numerous that one begins to wonder how any plant can survive with so many toxins collected in and on the soil. The answer is, of course, that normally such toxic products do not




Figure 15. Overstory-understory relationships vary with different soils.

accumulate but are broken down by soil microorganisms. Good descriptions of the breakdown processes are found in Harborne (1964) (fig. 16) and Schubert (1965). If a serious problem of noncompetitive antagonism appears after a study of interspecific relationships, then a study of possible chemical affects is called for, including a study of soil microorganism and the various physical and chemical processes that influence the microbial activity.

IX. Effect of Grazing.

Effect of grazing on herbaceous plants has been reviewed by Ellison (1960) and Jameson (1963). In the case of shrubs there is a need for studies of the effect of grazing (fig. 17). Some work along this line is being done at Tonto Springs, but this work will need to be supplemented with additional work.

X. Higher Animals, Forage Consumption, and Supplemental Feed

Present work at Wild Bill and Tonto Springs is designed to show the relationship between consumption of digestible forage and animal weight gains.

At the Wild Bill study area in the ponderosa pine type the relationships are very close to the expected. At the Tonto Springs study area in the chaparral type, however, the animal output in the brush pastures in the winter is much too high for the grass forage inputs. The problem is that we had identified shrub live oak as a loss from the stationary processor when actually it is being used as an input to the moveable processor. During the winter, from 55 to 65 percent




Figure 16. Normal process of litter breakdown.



Figure 17. Regrowth on grazed shrub live oak.

of the animal product output does not come from the grass under these conditions, and must be coming from the brush (table 2).

Two refinements are needed in these studies:

(1) The digestibility should be expressed as digestible energy; a high speed method of energy determination is needed for this analysis.

(2) Automatic weighing of animals is needed to reduce or eliminate weight losses due to handling.

So far all of our work has been done with yearling beef animals. We have no documented knowledge of cow-calf operations on these ranges. This deficiency of knowledge is most severe in wintering pregnant cows on chaparral ranges, least severe with cows and calves on pine and juniper summer ranges, and intermediate in wintering pregnant cows on juniper ranges.

The work at Tonto Springs has shown that shrub live oak is a major component of the cattle diet. We have no knowledge of the effect of very high percentages of shrub live oak intake on rumen ecology, cattle weight gains, and pregnancies. Calf crops in the chaparral are traditionally low; the cause is undetermined but it may be due to abortion inducing substances in the diet. Studies should be initiated to cover these aspects.

A beginning study on rumen ecology can begin with microorganism populations in in vitro cultures and microchemical and bioassay tests of possible toxins in the feed. Effects of feeds on abortion in ruminants likely will require more time, money, and effort.

Table 2. Grass and forb consumption (pounds per
pasture) at Tonto Springs

| Type of pastures | 1964 | 1965 |
|---------------------------------------|--------|--------|
| Native | 6,800 | 9,400 |
| Rootplowed | 14,746 | 25,974 |
| Native as % of plowed | 46% | 36% |
| Deficit in native forb consumption | 54% | 64% |

No severe nutritional deficiencies are apparent in pine summer ranges or chaparral yearlong ranges. There are likely to be protein deficiencies on pinyon-juniper winter ranges and in such areas supplementing with a protein supplement is a common practice. If further studies show that high levels of shrub live oak in the diet are detrimental there may be opportunities for correcting the problem with selected supplemental feed. During the period of time covered by this analysis, however, such studies will not progress beyond the preliminary test tube stage because we do not know yet whether or not a problem exists.

We suspect that supplemental feeds may change the value, and perhaps the palatability of some forages, especially browse, and we plan to investigate this. Economic feedback can be especially effective on the stationary processor where there is presently a loss that could be easily converted to an output. Chaparral species and tobosa grass are two examples. By applying proper management techniques much of such potential loss can be utilized. The chaparral is a particularly urgent problem since there is a vast area of essentially untapped forage.

XI. The Economic Processor

Ranch operations in the pinyon-juniper type have been described (Cotner 1963) and the summer-only ranch operations in the ponderosa pine type are relatively simple. Some work on ranch operations in these types has been done and is being done in connection with the Beaver Creek Project. In the chaparral, however, we need knowledge of how the ranch units operate and what their problems are. Such a study would undoubtedly point out other avenues for research.

XII. Brush Control and Reseeding

Brush control and grass reseeding in the problem area is handled by a team of Agricultural Research Service scientists located at Flagstaff. Good recommendations for reseeding in the chaparral and ponderosa pine types are available. Current ARS research is emphasizing the pinyon-juniper type and they are having considerable success with Luna pubescent wheatgrass and other species. In addition, Project 1704 is conducting research in browse plantings.

One possibility that is not presently being emphasized is planting of Ceanothus greggii (fig. 18). This species is a high producing, palatable browse plant which is not naturally present in abundance. Other chaparral browse plots are either abundant or unpalatable or both, but a greater amount of Ceanothus would be helpful.

In brush control there are many problems to be solved; in most cases other agencies and other uses are involved.

The ponderosa pine type represents the potentially highest producing area. Much of the ponderosa pine type is greatly overpopulated with pine trees, even from a forestry standpoint. At Wild Bill and Taylor Woods (Studies 14 and 15) the relationship between trees and herbage production is being studied. Some soils obviously do not have the same relationship between understory and overstory as the Taylor Woods-Wild Bill area (fig. 19). Manzanita (fig. 20), for example, is characterized by very low understory production. Since detailed studies of



Figure 18. Ceanothus greggii: a high producing and palatable
but relative scarce browse species of the chaparral.



Figure 19. Some soils such as this decomposed granite support
little understory vegetation.



Figure 20. Manzanita stands are characterized by lack of understory vegetation.

all soils would be impractical the first step should be a survey type study to determine which soils are problem soils in this regard. This and studies on litter breakdown discussed earlier should lead to future studies with direct management applications.

In the pinyon-juniper type the problem soils with regard to brush control have been identified. There are suitable mechanical brush control treatments available and also an accepted burning procedure for individual trees. A study on broadcast burning has been outlined (see section on program). ARS is working on chemical methods of control, but as yet these methods are rather expensive.

In the chaparral brush control studies are also underway by ARS. Further work in this area should await evaluation of ARS chemical control work, the Tonto Springs grazing study, and the prescribed burning program being conducted on the Tonto National Forest. At present it appears that areas with basalt soil in the chaparral can be managed for grazing without brush control; granitic outwash areas can be rootplowed.

XIII Fertilizers

A very little work, except for one study by Lavin (1967) has been done on range fertilization in the problem area. Economic use of macro nutrients will undoubtedly be restricted to the higher rainfall areas; these rainfall restrictions should be identified. Use of micro nutrients, however, may be feasible over a much wider range of conditions. A

comprehensive study of micro nutrients should be undertaken. For maximum efficiency such a study should begin with greenhouse pot tests. This will of course require a greenhouse. At the completion of the pot test phase field plots should be established.

XIV. The Range System

Recent developments in mathematics and computers have led biologists to attempt simulation of ecological systems (e.g. Garfinkel 1962; Neel and Olson 1962; Odum 1960; Watt 1959). Results so far have been very encouraging, and there is real hope for progress in such a concept. It is perhaps noteworthy that biological systems analysis has been developed by medical doctors, and other biologists rather than by mathematicians. It has been pointed out earlier in this analysis that even our crude system concept has pointed out errors in our thoughts of the range system. A good starting point for such research would be in a better documented type such as the ponderosa pine; this should provide a framework for the chaparral type where less information is available.

PRESENT PROGRAM

Specifically, the present program includes the following:

Study No. 8

TECHNIQUES OF BROADCAST BURNING IN DENSE

STANDS OF PINYON-JUNIPER TREES

Objective

Develop suitable methods of broadcast burning pinyon-juniper stands using chemical desiccants as an aid.

Status

Preliminary individual tree testing has been completed and the method shows promise. Large scale testing can be initiated as soon as an interested land management agency can be induced to treat a selected study area for an approximate cost of \$2,000.

Leader.--Jameson

Comments.--This is a fire use study and could best be handled by fire research. A pilot study was initiated in 1967 on the Beaver Creek Watersheds.

PRODUCTIVE POTENTIAL OF SITES IN THE PINYON-JUNIPER TYPE

Objectives

The overall objective of the study will be to determine productivity of various sites of the pinyon-juniper type expressed in grass production. This is to be accomplished in three parts; the first two parts have been completed.

1. A survey of the common boundaries of the tertiary volcanics and other geologic formations within the pinyon-juniper type to establish paired study plots at these boundary areas.
2. A comparison of the fertility and physical properties of soils developed from tertiary volcanics to determine the comparability of the volcanic soils at different locations.
3. To compare the productivity of selected pairs of plots selected on the basis of knowledge gained under objectives 1 and 2.

Status

All plots have been fenced and trees removed. Weeds were sprayed in 1965 and 1966, herbage samples were collected in 1964 and 1965. Herbage sampling will continue in 1967, 1969, 1971, and 1974. Weed spraying will be repeated when needed.

Leader.--Jameson. This study could be handled equally as well by Pond, but will not be transferred because of Pond's other workload.

AUTECOLOGY AND PLANT COMPETITION IN THE PINYON-JUNIPER TYPE

Objectives

1. To gain information on the seasonal patterns of growth and development of some important forage plants of the pinyon-juniper type which can be used in developing management systems.
2. To determine the differences in establishment and production of grasses when released from competition of herbaceous weeds as compared to nonrelease.
3. To determine the presence and investigate the importance of germination inhibitors in weeds, trees, grasses, plant litter, and soils of the pinyon-juniper type.
4. To determine the nature of the effect of pinyon and juniper trees on grasses, and the effect of grasses on trees.
5. To determine the regularity and abundance of seed production and requirements for seed germination of important native forage plants of the type.
6. To determine the effects of unstable soils on plant establishment and growth.
7. To determine the climatic patterns of the type so that realistic management goals may be established within the various patterns.

Status

Objectives 1, 2, 4 have been completed and reports published. Work is continuing on objectives 3 and 6. Objective 7 has been covered by work of other agencies. No work has been done on objective 5.

Leader.--Jamerson

PONDEROSA PINE OVERSTORY EFFECT ON RANGE PRODUCTION

Objectives

The objectives of this research will be to study the overstory-understory relationships as follows:

PHASE I (Burro Mountain)

To determine the effect that different cutting practices and slash disposal methods have on the composition, cover and production of the herbaceous and shrubby vegetation of the mixed conifer type.

PHASE II (Taylor Woods)

1. To determine the effects of tree overstories on forage production.
2. To determine the effects of overstory treatment practices on forage production.

Status

Phase I is presently not yielding useful information and has been closed.

Phase II. Pretreatment measurements and measurements 2 and 4 years after treatment have been completed.

EFFECT OF PONDEROSA PINE OVERSTORY ON FORAGE

AND BEEF PRODUCTION

Objectives

The objective of this study will be to compare animal production on different degrees of conversion from pine overstory to grass:

1. To determine the effects of tree overstories on forage production.
 - a. Quantity and composition of livestock and game forage
 - b. Quality
 - (1) Palatability and grazing and browsing patterns
 - (2) Production of digestible nutrients
2. To determine the effects of various tree overstories on the value of forage and tree production.

Status

Vegetation and animal production data have been collected for 4 years on the untreated area and 2 years on the treated areas. The study should be continued 3 to 6 years under the present objectives.

Leader.--Pearson. This study and No. 104 both have aspects which are best suited for work by Pearson, but they also have aspects which are best suited for work by Pond.

IDENTIFICATION AND ECOLOGICAL IMPORTANCE OF NATURALLY
OCCURRING GROWTH INHIBITORS OF SOUTHWESTERN
FOREST AND RANGE PLANTS

Objectives

1. To determine the presence of plant phenolic compounds which persist following normal decomposition processes.
2. To identify, if possible, the persistent chemicals.
3. To determine factors which can be manipulated that break down the persistent chemicals.

Status

Procedures have been developed which have allowed identification of the active inhibitors in juniper as isomes of chlorogenic acid. These acids are oxidized by the enzyme tyrosinase which may prove to be the clue to more rapid breakdown.

Leader.--Jameson

COMPARISON OF CATTLE GAINS AND FORAGE PRODUCTION

FROM CHAPARRAL IN NATURAL AND ROOTFLOWED

AND SEEDED CONDITION

Objectives

The major objectives of this study are to: (1) Compare year-long animal production from grazing on (a) rootplowed and seeded chaparral and (b) on native chaparral and (2) compare animal production from grazing rootplowed and seeded pastures in summer and native chaparral in winter with animal production from pastures in native and seeded condition grazed year long.

Status

The third year of data, animal gain, and forage production is almost complete. Grazing pressure has been increased on all pastures and now approaches the desired level (40%) on rootplowed and seeded pastures. Proper use of native pastures, especially the browse, is difficult to obtain since little is known concerning proper browse utilization. However, cattle gains from both types have been comparable. Until some difference in animal gain, seasonal or otherwise, becomes apparent between types, additional animals must be added to chaparral pastures each year. So far, utilization of brush species, especially shrub live oak, has been negligible as measured by our techniques.

Leader.--Pond

GROWTH, DEVELOPMENT AND NUTRITIVE VALUE

OF SOME CHAPARRAL SHRUBS

Objectives

The primary objective of this study is to gather information concerning the growth, development and nutritive value of several important shrubs of the chaparral type in Arizona. In so doing, the following specific questions will be answered:

1. When does growth begin and how long do the shrubs continue to grow during any one year?
2. What factors (rainfall, soil moisture, temperature) appear to regulate growth?
3. What is the crude protein, moisture and crude fiber content of current annual production at various stages of the plant's seasonal development?

All data

Status

All data has been collected and summarized. A manuscript is in preparation and will close this study.

Leader.--Pond

CHEMICAL AND BOTANICAL COMPOSITION OF THE DIET OF FISTULATED
STEERS GRAZING IN THE ARIZONA CHAPARRAL

Objectives

The objectives of this chaparral grazing study are (1) to estimate the botanical composition of the diet of cattle, (2) to estimate digestibility of the cattle diet, (3) to compare the nutrient content of the cattle diet with the main forage producing species, and (4) to estimate forage intake of the cattle.

Status

Cattle diet and forage species samples were periodically collected from December 1965 through October 1966. These collections should be continued for 2 more years.

Leaders.--Pearson and Pond

STAFFING

The project is presently underfinanced for the existing staffing, and additional staffing must be accompanied by increased financing.

Addition of 2 scientists would bring the project up to the strength of the 10-year program, but 3 would be better in view of the diversity of the problems. Additional scientists would also necessitate additional support personnel (table 3). A proposed organization is as follows:

Project Leader

Provides project supervision and guidance. Conducts research in specialty field.

Animal Physiologist

Conducts digestibility studies on range browse and forage species, determines impact of "undesirable" diets such as shrub live oak on animal weights and reproduction. Most of this scientist's time and his basic research will be in the chaparral, although he will supervise digestion studies at Wild Bill and Beaver Creek which are by now mostly routine.

Range Scientist or Agronomist

Conducts agronomic type experiments where treatments are applied and vegetation response is measured in the field. Treatments are of field plot or pilot plant size. Treatments may include mechanical, chemical, or fire control of brush and addition of micro nutrients, or perhaps macro nutrients. Work will be about evenly divided among the pine, pinyon-juniper, and chaparral type.

Biosystems Analyst

Analyzes relationships between components of the range system. These components include environment, soil, plant and microorganism community, domestic animals, and ranch operations. Research techniques will resemble linear programming of economists and systems analysis of electronic specialists. This is an especially promising area of research, but recruiting a systems analyst with a good knowledge of biology will be a problem. Most need is in the chaparral type.

Microflora Ecologist

Studies impact of plant toxins which accumulate under conifers and in heavy clay soils. Develops methods of promoting breakdown of toxins. Most need is in the pinyon-juniper type, secondary need is in the ponderosa pine type.

Soil Scientist

Studies relationship of grazing practices to soil erosion. Particular need is in the chaparral type where management of the range for the most abundant species (shrub live oak) is suspected of creating an erosion hazard. Management for the grass species, on the other hand, utilizes only about 5 percent of the potential of the type.

Experimental Area Superintendent

General supervision of larger experimental areas, including routine field data collection as specified by the scientists.

Technician (2)

Collection of field data.

Laboratory Technician (2)

Conduct routine laboratory analyses, particularly in artificial rumen digestion studies and soil microflora studies. An assistant will be needed during the summer months.

Data Processing Technician

Routine data processing, including key punch and automatic data processing equipment.

Greenhouseman

Provides routine maintenance of greenhouse and growth chamber studies.

Table 3. Staffing (man-years) and budget

| Type of personnel | Existing | Needed for existing scientists | Needed with 3 additional scientists |
|----------------------------------|----------|--------------------------------|-------------------------------------|
| Scientists | 3 | 3 | 6 |
| Experimental area superintendent | 0 | 1 | 1 |
| Field technicians, full time | 3 | 1 | 3 |
| Greenhouseman | 0 | 0 | 1 |
| Laboratory technicians | 0 | 1 | 2 |
| Summer field assistants | 0 | | 2 (6x4 mos) |
| Statistical clerk | 0 | 1/2 | 1 |
| Clerk stenographers | 1/2 | 1/2 | 1 |
| Approximate budget | \$78,000 | \$130,000 | \$280,000 |

ADDITIONAL STUDIES PROPOSED WITH PRESENT STAFFING

1. Treatments for promoting sprouts of chaparral browse species. (Proposed for Pond)
2. Microorganism population with single and mixed in vitro diets of chaparral species. (Proposed for Pearson)
3. Ecology of snakeweed. Only preliminary investigation can be made without greenhouse facilities and with present staff. (Proposed for Jameson)
4. Establishment of Ceanothus greggii. In this study only preliminary investigations can be made without a greenhouse. (Proposed for Pond with some phases handled by Springfield)
5. Effect of shrubs on phenology and distribution of grasses. (Proposed for Pond)

Ecology + Phenology
of shrubs of grasses.

ADDITIONAL STUDIES WHICH COULD BE UNDERTAKEN WITH
ADEQUATE GREENHOUSE FACILITIES

1. Ecology of Gutierrezia (Expansion of proposed study)
2. Establishment of Ceanothus greggii. (Expansion of proposed study)
3. Micro nutrient deficiencies in soils.

ADDITIONAL STUDIES WHICH COULD BE UNDERTAKEN

WITH ADDITIONAL STAFFING

1. Relationship between grazing and erosion in the chaparral. This study could be done by the watershed research group and should at least be done in cooperation with that group.
2. Basic rumen ecology in the chaparral type.
3. Survey of soil microorganism populations. Alternatively, this could be handled under cooperative agreement.
4. Fertilization trials in the pine and chaparral types. Community ecology on alluvian, basalt, and residual igneous and metamorphic sites of the Arizona chaparral.
5. Range Systems Analysis. Nowhere in the range management field is there a study of systems analysis such as there is in insect population dynamics and predator-prey relationships. A system analysis investigation back up by a staff of biologists should yield good results.

COOPERATIVE STUDIES

Cooperative studies needed in conjunction with either the abbreviated or expanded program include:

(1) An economic analysis of ranch operation in the Arizona chaparral.

(2) A survey of soil microorganism populations on selected range soils.

PROJECT PUBLICATIONS

PUBLICATIONS, ADDRESSES, AND MANUSCRIPTS

Jameson, Donald A.

1959. Use of fire in control of juniper in Arizona. Paper presented to Western Weed Control Conf. 1959.

Cotner, M. L., and D. A. Jameson

1959. Costs of juniper control: bulldozing vs. burning individual trees. Rocky Mountain Forest and Range Expt. Sta. Paper 43.

Jameson, Donald A.

1961. Heat and desiccation resistance of tissue of important trees and grasses of the pinyon-juniper type. Bot. Gaz. 122: 174-179.

-
1961. Growth inhibitors in native plants of northern Arizona. Rocky Mountain Forest and Range Expt. Sta. Res. Note 61, 2 pp.

_____, J. A. Williams, and E. W. Wilton

1962. Vegetation and soils of Fishtail Mesa, Arizona. Ecology 43: 403-410.

Jameson, Donald A.

1962. Effects of burning on a galleta-black grama range invaded by juniper. Ecology 43: 760-763.

-
1963. Evaluation of the responses of individual plants to grazing. In Range Res. Methods. A symposium. U.S. Dept. Agr. Forest Serv. Misc. Pub. 940. 109-116.

-
1963. Physiology of defoliation. Paper presented to joint meeting of American Society of Agronomy and American Society of Range Management. See: Agron. Abstr. 17-21: 105.

-
1963. Responses of individual plants to harvesting. Botan. Rev. 29: 532-594.

_____, and T. N. Johnsen, Jr.

1964. Ecology and control of alligator juniper. Weeds 12: 140-142.

Arnold, J. F., D. A. Jameson, and E. H. Reid

1964. The pinyon-juniper type of Arizona: effects of grazing, fire, and tree control. U.S. Dept. Agr. Prod. Res. Rpt. 84, 28 pp.

Jameson, Donald A.

1964. Effects of defoliation on forage plant physiology. Amer. Soc. Agron. Spec. Publ. 5, 67-80.

1964. Effect of grazing on plants. Paper presented to Arizona Section of the American Society of Range Management.

1965. Arrangement and growth of pinyon and one-seed juniper trees. Plateau 37: 121-127.

_____, and E. H. Reid

1965. The pinyon-juniper type of Arizona. Jour. Range Mangt. 18: 152-153.

1965. Phenology of grasses of the northern Arizona pinyon-juniper type. U.S. Forest Serv. Res. Note RM-47, 8 pp.

1965. Reduction of understory grass growth by chemicals in litter of Juniperus. Paper presented to joint meeting of Ecological Society of America, American Society of Plant Physiologists, and Botanical Society of America.

1966. Competition in a blue grama-broom snakeweed-actinea community and responses to selective herbicides. Jour. Range Mangt.

1966. Pinyon-juniper litter reduces growth of blue grama. Jour. Range Mangt.

1966. Diurnal and seasonal fluctuations in moisture content of pinyon and juniper. U.S. Forest Serv. Res. Note In press.

1966. Juniper control by individual tree burning. U.S. Forest Serv. Res. Note RM-71, 4 pp.

Jameson, Donald A.

1966. A growth function for the relationship of tree overstory and herbaceous understory. Jour. Range Mangt. In Press.

Pond, Floyd W., and Dwight R. Cable.

1960. Effect of heat treatment on sprout production of some shrubs of the chaparral in central Arizona. Jour. Range Mangt. 13(6): 313-317.

Pond, Floyd W.

1964. Response of grasses, forbs, and halfshrubs to chemical control of chaparral in central Arizona. Jour. Range Mangt. 17(4): 200-203.

1964. Toward more successful rootplowing. Arizona Cattlelog 21(3): 2 pp.

 , D. T. Lillie, and H. R. Holbo

1965. Shrub live oak control by rootplowing. U.S. Forest Serv. Rocky Mountain Forest and Range Expt. Sta. Res. Note 38, 2 pp.

Pond, Floyd W.

1961. Effect of three intensities of clipping on the density and production of meadow vegetation. Jour. Range Mangt. 14(1): 34-38.

1961. Basal cover and production of weeping lovegrass under varying amounts of shrub live oak crown cover. Jour. Range Mangt. 14(6): 2 pp.

1961. Mechanical control of Arizona chaparral and some results of brush clearing. "Modern Techniques in Water Management," Proceedings, Fifth Annual Arizona Watershed Symposium, 3 pp.

1962. Shrub live oak limits production of weeping lovegrass. Arizona Cattlelog. 13(7): 61.

 , and Dwight R. Cable

1962. Recovery of vegetation following wildfire on a chaparral area in Arizona. Rocky Mountain Forest and Range Expt. Sta. Res. Note 72, 4 pp.

Pearson, Henry A.

1964. Beef--trees--or both. Arizona Cattlelog. 8-9.

Pearson, Henry A.

1964. Studies of forage digestibilities under ponderosa pine stands. Soc. Amer. Foresters Proc. 71-73.

1965. Low-cost temperature water bath. Jour. Range Mangt. 18(3): 149-150.

1965. Rumen organisms in white-tailed deer from South Texas. Jour. Wildlife Mangt. 20(3): 493-496.

UNPUBLISHED MANUSCRIPTS

Clary, Warren P., and Henry A. Pearson

Preference ratings of forage species in northern Arizona.

_____, Peter F. Ffolliott, and Donald A. Jameson

Effect of different forest floor layers on herbage production.

Pearson, Henry A.

Comparison of paired plot techniques used in determining forage utilization.

A consideration of inherent animal-grazing abilities in range grazing studies.

Digestibility of range cattle diet related to digestibility of component species.

Effect of delays in inoculum collection on artificial rumen digestibilities.

Phenology of two important range grasses of the Northern Arizona ponderosa pine type.

_____, and Donald A. Jameson

Relationship between timber and cattle production on ponderosa pine range--the Wild Bill Study.

Pond, Floyd W.

Declining grass density and production on ungrazed chaparral areas converted to grass.

_____, and Arthur R. Tiedemann

Viability of grass seed stored for long periods of time without temperature or humidity control.

OTHER LITERATURE

The literature included here has not all been cited in this project analysis. It has not been brought together before, however, and such a listing should be worthwhile. An examination of this list shows an abundance of background information in the pine and pinyon-juniper types but little in the chaparral.

References not found in the main alphabetical listing will be found in a supplemental list beginning on page 112.

- Adams, John E., and R. J. Hanks.
1964. Evaporation from soil shrinkage cracks. Soil Sci. Soc. Amer. Proc. 28(2): 281-284.
- Aldon, Earl F.
1964. Ground-cover changes in relation to runoff and erosion in West-central New Mexico. U.S. Forest Serv., Rocky Mountain Forest and Range Expt. Sta., Res. Note RM-34, 4 pp. Jour. Range Mangt. 18(6): 351.
- Allred, B. W.
1949. Distribution and control of several woody plants in Oklahoma and Texas. Jour. Range Mangt. 2: 17-29.
- Altschuler, E. S., and E. J. Dwornik
1963. Transformation of montmorillonite to kaolinite during weathering. Science 141: 148-152.
- Anderson, D., and others.
1953. Reseeding desert grassland ranges in southern Arizona. Ariz. Agr. Expt. Sta. Bul. 249, 32 pp., illus.
- Anderson, Henry W.
1947. Soil freezing and thawing as related to some vegetation, climatic, and soil variables. Jour. Forestry 45: 94-101.
- Armer, W.
1952. Larkspur, loco, and lupine. Ariz. Cattlelog, pp. 14-16. July.
- Arnold, J. F.
1942. Forage consumption and preferences of experimentally fed Arizona and antelope jackrabbits. Ariz. Agr. Expt. Sta. Tech. Bul. 98: 51-86.
- _____
1950. Changes in ponderosa pine-bunchgrass ranges in northern Arizona resulting from pine regeneration and grazing. Jour. Forestry 48: 118-126.
- _____
1953. Effect of heavy selection logging on the herbaceous vegetation in a ponderosa pine forest in northern Arizona. Jour. Forestry 51: 101-105.
- _____
1955. Juniper control. Ariz. Cattlelog 10(9): 44-50

Arnold, J. F.

1955. Juniper control as a range-improvement practice in central Arizona. Ranch Day New Mex. Expt. Sta. and Rocky Mountain Forest and Range Expt. Sta., Fifteenth Ann. Ranch Day. (Mimeo)

1955. Plant life form classification and its use in evaluating range conditions and trend. Jour. Range Mangt. 8(4): 176-184.

1964. Zonation of understory vegetation around a juniper tree. Jour. Range Mangt. 17: 41-42.

_____, D. A. Jameson, and E. H. Reid

1964. The pinyon-juniper type of Arizona: Effects of grazing, fire, and tree control. U.S. Dept. Agr. Forest Serv. Prod. Res. Rpt. No. 84, 28 pp.

Arnold, J. F., and W. L. Schroeder

1955. Juniper control increases forage production on the Fort Apache Indian Reservation. Rocky Mountain Forest and Range Expt. Sta. Sta. Paper 18, 35 pp., illus. (Mimeo)

Babbitt, J.

1950. Range problems of a northern Arizona ranch. Ariz. Cattlelog 5(11): 3-6.

Bailey, K. P.

1963. Influence of soil strength on growth of roots. Soil Sci. 96: 175-180.

Baker, F. S.

1944. Mountain climate of the western United States. Ecol. Monogr. 14: 223-254.

Barnes, O. K., D. Anderson, and A. Heerwagen

1958. Pitting for range improvement in the Great Plains and the Southwest desert regions. U.S. Dept. Agr. Prod. Res. Rpt. 23, 17 pp., illus.

Barr, George V.

1955. A comparison of the survival over a twenty-year period of several native Arizona grasses. M.S. Thesis, Univ. of Ariz.

Barr, George W.

1956. Recovering rainfall. Ariz. Watershed Program (Ariz. State Land Dept., Salt River Valley Water Users Assn., Univ. of Ariz., cooperating), 33 + 218 pp., illus.

Basnett, P., and others.

1940. Report on the interagency study of the cedar problem on the Edwards Plateau and the adjacent areas in the central and south-central Texas. Unpubl. 4 pp.

Bates, C. G.

1923. Physiological requirements of Rocky Mountain trees. Jour. Agr. Res. 24: 97.

Bell, H. M., and E. J. Dyksterhuis

1943. Fighting the mesquite and cedar invasion on Texas ranges. Soil Conserv. 9: 111-114.

Benedict, H. M.

1940. Effect of day length and temperature on the flowering and growth of four species of grasses. Jour. Agr. Res. 61: 661-672.

- _____. 1941. Growth of some range grasses in reduced light. Botan. Gaz. 102: 582-589.

_____, and G. B. Brown

1944. The growth and carbohydrate responses of Agropyron smithii and Bouteloua gracilis to changes in nitrogen supply. Plant Physiol. 19: 481-494.

Berndt, Herbert W., and Robert D. Gibbons

1958. Root distribution of some native trees and understory plants growing on three sites within ponderosa pine watersheds in Colorado. U.S. Forest Serv., Rocky Mountain Forest and Range Expt. Sta. Sta. Paper 37, 14 pp.

Berry, W. E.

1948. Water spreading for soil and moisture conservation in the Southwest. U.S. Dept. Int. Bur. of Indian Affairs, 18 pp., illus.

_____, and D. Anderson

1943. The effect of surface mulches on water conservation and forage production in some semi-desert grassland soils. Jour. Amer. Soc. Agron. 35: 393-400.

Beveridge, W., and C. Ames

1956. Does juniper control pay? Results obtained on the West Bear allotment, Prescott National Forest. Report to 1956 summer meeting of Ariz. Sec., Amer. Soc. Range Mangt., 5 pp.

Billings, W. D.

1951. Vegetational zonation in the Great Basin of western North America. C. R. du Colloque sur les bases ecologiques de la regeneration de la vegetation des zones arides. U. I. S. B. 101-122. Paris.

-
1954. Temperature inversions in the piñon-juniper zone of Nevada Mountain Range. Butler Univ. Botan. Studies XI: 112-118.

Börner, H.

1960. Liberation of organic substances from higher plants and their role in the soil sickness problem. Botan. Rev. 26: 393-424.

Bostick, V. B.

1947. Principles for judging condition and trend of southwestern woodland ranges. U.S. Forest Serv., Region 3, 50 pp.

Bradshaw, K. E., and J. L. Reveal.

1943. Tree classifications for Pinus monophylla and Juniperus utahensis. Jour. Forestry 41: 100-104.

Bray, W. L.

1940. The timber of the Edwards Plateau of Texas; its relation to climate, water supply, and soil. U.S. Dept. Agr. Bur. of Forestry Bul. 49, 30 pp.

Brown, H. Ray

1943. Growth and seed yields of native prairie plants in various habitats of the mixed prairie. Trans. Kans. Acad. Sci. 46: 87-99.

Bryson, Reid A.

1957. The annual march of precipitation in Arizona, New Mexico, and northwestern Mexico. Univ. of Ariz., Inst. of Atmos. Physics Tech. Rpts. on the Meteorology and Climatology of Arid Regions No. 6, 24 pp.

Buchanon, Hoyle.

1961. The plant ecology of Bryce Canyon National Park. Diss. Abstr. 21: 2445.

Cable, D. R.

1957. Chemical control of chaparral shrubs in central Arizona.
Jour. Forestry 55: 899-903.

1957. Recovery of chaparral following burning and seeding in central Arizona. U.S. Forest Serv., Rocky Mountain Forest and Range Expt. Sta. Res. Note 28, 6 pp., illus.

_____, and J. W. Bohning

1959. Changes in grazing use and herbage moisture content of these exotic lovegrasses and some native grasses.
Jour. Range Mangt. 12: 200-203.

_____, and F. H. Tschirley

1961. Responses of native and introduced grasses following aerial spraying of velvet mesquite in southern Arizona. Jour. Range Mangt. 14: 155-159.

Campbell, R. S.

1943. Progress in utilization standards for western ranges.
Wash. Acad. Sci. Jour. 33: 161-169.

1946. Determination of grazing values of native vegetation on southern pine forest ranges. Ecology 27: 195-204.

1955. Vegetational changes in the cut over longleaf pine-slash pine area of the Gulf Coast. Ecology 36: 29.

_____, and E. H. Bomberger

1943. The occurrence of *Gutierrezia sarothrae* on *Bouteloua eripoda* ranges in southern New Mexico. Ecology 15: 49-61.

Canfield, R. H.

1957. Reproduction and life span of some perennial grasses of southern Arizona. Jour. Range Mangt. 10: 199-203.

Carter, W. T., and V. L. Cory

1932. Soils of trans-Pecos Texas and some of their vegetative relations. Tex. Acad. Sci. Trans. and Proc. 15: 19-32.

Cassady, J. T.

1937. Chamiza browse on southwestern ranges, and ways to increase it. Southwestern Forest and Range Expt. Sta. Res. Note 23, 5 pp.

Cassady, J. T.

1937. Scarification of hardcoated grass seeds to increase germination. Southwestern Forest and Range Expt. Sta. Res. Note 13, 2 pp.

Cassidy, H. O.

1937. How cattle may use cut-over ponderosa pine-bunchgrass with minimum injury to reproduction. Southwestern Forest and Range Expt. Sta. Res. Note 32, 15 pp.

1937. How to handle sheep on cut-over ponderosa pine-bunchgrass ranges with minimum injury to reproduction. Southwestern Forest and Range Expt. Sta. Res. Note 16, 2 pp.

Catlin, G. N.

1925. Composition of Arizona forages, with comparative data. Ariz. Agr. Expt. Sta. Bul. 113; 155-173.

Chapline, W. R.

1919. Production of goats on far western ranges. U.S. Dept. Agr. Bul. 749, 35 pp., illus.

_____, and M. W. Talbot.

1926. The use of salt in range management. U.S. Dept. Agr. Dept. Cir. 379, 32 pp., illus.

Christensen, Earl M.

1964. Changes in composition of a Bromus tectorum--Sporobolus cryptandrus--Aristida longiseta community following fire. Utah Acad. Sci. Arts and Letters Proc., Part I, 41: 53-57. Jour Range Mangt. 18(6): 350.

Colton, Harold S.

1958. Precipitation about the San Francisco Peaks. Museum of Northern Arizona Tech. Ser. 2.

Cook, C. Wayne, L. A. Stoddart, and L. E. Harris

1959. The chemical content in various portions of the current growth of salt-desert shrubs and grasses during winter. Ecology 40: 644-651.

Cooper, C. F.

1960. Production of native and introduced grasses in the ponderosa pine region of Arizona. Jour. Range Mangt. 13: 214-215.

Cooper, C. F.

1961. Controlled burning and watershed condition in the White Mountains of Arizona. Jour. Forestry 59(6): 438-442.

Cooperrider, Charles K., and Cassidy, Hugh O.

1941. Cattle grazing on cutover timberlands in relation to regeneration of ponderosa pine forests of the Southwest. Files, Rocky Mountain Forest and Range Expt. Sta., Flagstaff, Ariz.

_____, and B. A. Hendricks

1940. The role of vegetation on steep mountain slopes. Southwestern Forest and Range Expt. Sta. Res. Note 85, 3 pp.

Corte, Arturo E.

1963. Particle sorting by repeated freezing and thawing. Science 142: 499-501.

Cory, V. L.

1956. Three junipers of western Texas. Rhodora 58: 183-187.

Cotner, Melvin L.

1963. Controlling pinyon-juniper. Ariz. Agr. Expt. Sta. Rpt. 210, 28 pp.

- _____, 1963. Optimum timing of long-term improvements. Jour. Farm Econ. 45: 732-748.

_____, and Donald A. Jameson

1959. Predicting costs of juniper control by bulldozing and individual-tree burning. U.S. Forest Serv., Rocky Mountain Forest and Range Expt. Sta. Sta. Paper 43, 14 pp., illus.

_____, and M. M. Kelso

1963. Northern Arizona beef production is affected by pinyon-juniper. Prog. Agr. in Ariz. 15 (2).

Cottle, H. J.

1931. Studies in the vegetation of southwestern Texas. Ecology 12: 105-155.

- _____, 1932. Vegetation on north and south slopes of mountains in southwestern Texas. Ecology 13: 121-134.

Coupland, Robert T.

1958. The effects of fluctuations in weather upon the grasslands of the Great Plains. Botan. Rev. 24: 273-318.

Crafts, E. C.

1937. Proper utilization of bunchgrass range. Southwestern Forest and Range Expt. Sta. Res. Note 12, 3 pp.

1938. Tentative range utilization standards, bluestem (Agropyron smithii). Southwestern Forest and Range Expt. Sta. Res. Note 44, 5 pp.

1938. Tentative range utilization standards--side-oats grama (Bouteloua curtipendula). Southwestern Forest and Range Expt. Sta. Res. Note 36, 4 pp., illus.

1938. Tentative range utilization standards, blue grama (Bouteloua gracilis). Southwestern Forest and Range Expt. Sta. Res. Note 32, 4 pp.

1944. Growth and utilization of forage on southwestern pine ranges. Files, Rocky Mountain Forest and Range Expt. Sta., Flagstaff, Ariz.

_____, and G. E. Glendening.

1942. How to graze blue grama on southwestern ranges. U.S. Dept. Agr. Leaflet 215, 8 pp., illus.

Culley, M. J.

1946. Factors affecting range calf crop. Ariz. Stockman, 30-37. Nov.

1947. It pays to be loss conscious. Cattleman, 3 pp. Dec.

Dahms, Walter G.

1961. Chemical control of brush in ponderosa pine forests of central Oregon. Pacific Northwest Forest and Range Expt. Sta. Res. Paper 39, 17 pp., illus.

Dall, Howard

1955. Kill your Bull pines the easy way. West. Livestock Jour. 33(18): 38-39.

Darrow, R. A.

1944. Arizona range resources and their utilization. I.
Cochise County. Ariz. Agr. Expt. Sta. Tech.
Bul. 103: 311-366.

Daubenmire, R. F.

1943. Vegetational zonation in the Rocky Mountains. Botan.
Rev. 9: 325-393.

_____, and Don C. Presso

1963. Studies on the decomposition rates of tree litter.
Ecology 44: 589-592.

Deaver, Chester F., and Horace S. Haskell

1955. Pinyon resources - Navajo-Hopi Indian reservations -
Arizona-Utah. Univ. Ariz. Press., 37 pp.

Dodge, Richard Archie

1963. Investigations into the ecological relationships of
ponderosa pine in southeast Arizona. Diss. Abstr.
24(3): 917-918.

Dortignac, E. J., and L. D. Love

1961. Infiltration studies on ponderosa pine ranges of
Colorado. U.S. Forest Serv., Rocky Mountain
Forest and Range Expt. Sta. Sta. Paper 59, 34 pp.

Driscoll, Richard Stark.

- Characteristics of some vegetation-soil units in
the juniper zone in central Oregon. Diss. Abstr.
22(10): 3357-3358. (See Jour. Range Mangt. 1962)

Dryness, C. T., and C. T. Youngberg

1957. Effect of logging and slash burning on soil structure.
Soil Sci. Soc. Amer. Proc. 21: 444-447.

Dunford, E. G.

1949. Relation of grazing to runoff and erosion on bunchgrass
ranges. U.S. Forest Serv., Rocky Mountain Forest
and Range Expt. Sta. Res. Notes 7, 2 pp.

Dwyer, Don D.

1958. Competition between forbs and grasses. Jour. Range
Mangt. 11: 115-118.

Dyksterhuis, E. J.

1957. The Savannah concept and its use. Ecology 38(3):
435-442.

Eastwood, Alice

1919. Early spring at the Grand Canon near El Tovar. Plant
World 22: 95.

Ehrenreich, John H.

1959. Releasing understory pine increased herbage production.
U.S. Forest Serv., Central States Forest Expt. Sta.
Sta. Note 139, 2 pp.

1960. Usable forage under pine stands? U.S. Forest Serv.,
Central States Forest Expt. Sta. Sta. Note 142,
2 pp.

_____, and John S. Crosby

1960. Herbage production is related to hardwood crown cover.
Jour. Forestry 58: 564-565.

Ellison, Lincoln

1960. Influence of grazing on plant succession of rangelands.
Botan. Rev. 26: 2-78.

Emerson, F. W.

1932. Tension zone between the grama grass and pinyon-juniper
associations in northeastern New Mexico. Ecology
13: 347-358.

Empire Machinery Company

1960. Rootplowing. 12 pp.

Erdman, James Allen

1962. Ecology of the pinyon-juniper woodland of Wetherill
Mesa, Mesa Verde National Park, Colorado. MS
Thesis, U. of Colo.

Evanari, Michael

1957. The physiological action and biological importance
of germination inhibitors. Symp. Soc. Expt. Biol.
11: 21-43.

Evans, L. T., and E. W. Russell

1959. Adsorption of humic and fulvic acids by clays. Jour.
Sci. 10: 119-132.

Fenner, R. L., R. V. Arnold, and C. C. Buck

1955. Area ignition for brush burning. Calif. Forest and
Range Expt. Sta. Tech. Paper No. 10, 10 pp.

Fitcher, Edson

1953. Control of jack rabbits and prairie dogs on range lands. Jour. Range Mangt. 6: 16-24.

Fletcher, J. E., and E. L. Beutner

1941. Erodability investigations on some soils of the upper Gila watershed. U.S. Dept. Agr. Tech. Bul. 794, 32 pp., illus.

Forsling, C. L.

1929. The utilization of browse forage as summer range for cattle in southwestern Utah. U.S. Dept. Agr. Cir. No. 62, 29 pp., illus.

Frazer, J. W., and J. L. Farrar

1953. Effect of shade on jack pine germination. Silv. Leaflet Forest Br. Can. 88: 3.

Freeman, B. N., and R. R. Humphrey

1956. The effects of nitrates and phosphates upon forage production of a southern Arizona desert grassland range. Jour. Range Mangt. 9: 176-180.

Frischknecht, Neil C.

1951. Seedling emergence and survival of range grasses in central Utah. Agron. Jour. 43: 177-182.

-
1963. Contrasting effects of big sagebrush and rubber rabbitbrush on production of crested wheatgrass. Jour. Range Mangt. 16: 70-74.

Fuller, W. H., S. Shannon, and P. S. Burgess

1955. Effect of burning on certain forest soils of northern Arizona. Forest Sci. 1: 44-50.

Fulps, J. L.

1942. Somatic chromosome complements in Bouteloua. Amer. Jour. Botan. 29: 45-55.

Gaines, E. M., R. S. Campbell, and J. J. Brasington

1954. Forage production on longleaf pine stands of southern Alabama. Ecology 35: 59-62.

_____, and E. S. Kotok

1954. Thinning ponderosa pine in the Southwest. Rocky Mountain Forest and Range Expt. Sta. Sta. Paper 17, 20 pp., illus.

Galizioli, S.

1958. Deer-carrying capacity of various sub-types within the chaparral habitat. Ariz. Game and Fish Dept., Wildl. Restor. Div. Prog. Rpt., Proj. W-78-R2.

Garb, Solomon

1961. Differential growth-inhibitors produced by plants. Botan. Rev. 27: 422-443.

Garfinkel, D.

1962. Digital computer simulation of ecological systems. Nature 194: 856-857.

Garrison, George A., and Robert S. Rummell

1951. First year effects of logging on ponderosa pine forest lands of Oregon and Washington. Jour. Forestry 49: 708-913.

Gary, Howard L., and Lowell R. Rich

1961. Nitrogen aids plant growth on Arizona soils derived from granite and diabase. Rocky Mountain Forest and Range Expt. Sta. Res. Note 57, 2 pp.

Gill, W. R.

1961. Mechanical impedance of plants by compact soils. Amer. Soc. Agr. Engin. Trans. ASAE 4(2): 238-242.

Glendening, George E.

1939. Comparison of the effects of various kinds of artificial litter upon the germination of grass seeds. Southwestern Forest and Range Expt. Sta. Res. Note 61, 4 pp.

1942. Germination and emergence of some native grasses in relation to litter cover and soil moisture. Jour. Amer. Soc. Agron. 34: 797-804.

1944. Some factors affecting cattle use of northern Arizona pine-bunchgrass ranges. Southwestern Forest and Range Expt. Sta. Res. Rpt. 6, 9 pp., illus.

1948. Noxious plant control in the Southwest. A problem analysis. Files of Rocky Mountain Forest and Range Expt. Sta., pp. 22-29.

1951. Thoughts on the juniper question. Ariz. Stockman.

- Glock, Waldo S., Sharlene R. Agerter, and David G. Smith
1961. Recent change in the pattern of tree growth in northern Arizona. *Science* 134: 1417-1418.
- Graham, F. W.
1937. Botanical studies in the Uinta Basin of Utah and Colorado. *Ann. Carnegie Mus.* 26: 1-432.
- Gray, James R.
1956. Southwestern cattle ranches; organization, costs and returns. *New Mex. Agr. Expt. Sta. Bul.* 403, 87 pp.
- Griffiths, D.
1901. Range improvement in Arizona. U.S. Dept. Agr., Bur. Plant Ind. Bul. 4, 31 pp., illus.

1904. Range investigations in Arizona. U.S. Dept. Agr., Bur. Plant Ind. Bul. 67, 62 pp., illus.
- Grunner, G.
1961. The role of toxic substances in the interrelationship between higher plants. *Soc. Expt. Biol. Symp.* 15: 219-228.
- Hahn, Glen Gardner
1941. Reports of cedar blight in 1940 and notes on its previous occurrence in nurseries. U.S. Bur. Plant Indus. Plant Dis. Rpt. 25: 186-190.
- Hall, Marion T.
1955. Comparison of juniper population on an Ozark glade and old fields. *Ann. Missouri Botan. Garden* 42: 171-194.
- Halloran, A. F., and F. N. Ares
1944. Record of alligator juniper (*Juniperus pachyphloea* Torr.) in the Jornada Experimental Range. New Mexico. *Amer. Midland Nat.* 32: 518.
- Halls, Lowell K.
1955. Grass production under dense longleaf-slash pine canopies. *Southern Forest Expt. Sta. Res. Note* 83. July.
- Hanson, H. C.
1924. A study on the vegetation of northeastern Arizona. *Univ. Neb. Studies* 24: 1-94.

Hanson, William R.

1954. Ecological investigation of the Prescott Study Area.
Ariz. Game and Fish Comm. Proj. W-71-R-1, Work
Plan 4, Job 2, 9 pp.

_____, and Clay Y. McCulloch

1955. Factors influencing mule deer on Arizona brushlands.
Trans. 20th North Amer. Wildl. Cong. 568-588.

Harborne, J. B.

1964. Biochemistry of phenolic compounds. Academic Press,
N. Y. 618 pp.

Harlan, Jack R.

- Blue grama types from west Texas and eastern New
Mexico. Jour. Range Mangt. 11: 84-87.

Harmon, Claude F.

1951. The march of the cedars. Ariz. Cattlelog 6: 54-63.

Hawley, F. M.

1937. Cedar growth to precipitation and runoff. Ecology
18: 398-405.

Haskell, H. S., and H. G. Reynolds

1947. Growth, developmental food requirements, and breeding
activity of the California jackrabbit. Jour.
Mammology 28: 129-136.

Hastings, J. R.

1959. Vegetation change and arroyo cutting in southeastern
Arizona. Jour. Ariz. Acad. Sci. 1: 60-67.

_____, and Raymond M. Turner

1965. The changing mile. Univ. Ariz. Press. 317 pp.

Heinonen, Reijo

1962. On characteristics of top soils overlying heavy
clays. Maataloustieteellinen Aikakauskirja 34(1):
26-33.

Hendricks, B. A.

1936. Vine-mesquite for erosion control on southwestern
ranges. U.S. Dept. Agr. Leaflet 114, 8 pp., illus.

31

1939. Graded trails as an aid in the distribution of cattle
on rough ranges. Southwestern Forest and Range
Expt. Sta. Res. Note 67, 8 pp., illus.

Hendricks, B. A.

1942. Effect of grass litter on infiltration of rainfall on granitic soils in a semi-desert shrub-grass area. Southwestern Forest and Range Expt. Sta. Res. Note 96, 3 pp.

_____, and J. M. Johnson

1944. Effects of fire on steep mountain slopes in central Arizona. Jour. Forestry 42: 568-571.

Herman, F. R.

1953. A growth record of Utah juniper in Arizona. Jour. Forestry 51: 200-201.

_____, 1956. Growth and phenological observations of Arizona junipers. Ecology 37: 193-195.

_____, 1958. Silvical characteristics of Rocky Mountain juniper. Rocky Mountain Forest and Range Expt. Sta. Paper No. 29, 20 pp.

_____, and B. R. Lexen

1939. Fuel wood volume tables for Rocky Mountain red cedar (Juniperus scopulorum Sarg.) U.S. Forest Serv., Southwestern Forest and Range Expt. Sta. Res. Note 68, 2 pp.

Hoffman, A. F.

1921. The pinyon-juniper land problem. II. Plan for handling the pinyon-juniper type. Jour. Forestry 19: 537-541.

Holt, G. A., and D. G. Wilson

1961. The effect of commercial fertilizers on forage production and utilization on a desert grassland site. Jour. Range Mangt. 14: 252-256.

Honnas, R. C., B. L. Branscomb, and R. R. Humphrey

1959. Effect of range fertilization on growth of three southern Arizona grasses. Jour. Range Mangt. 12: 88-91.

Howell, Joseph, Jr.

1937. The relation of crown diameter to cubic volume of one-seed juniper. Jour. Forestry 35: 829-831.

Howell, Joseph, Jr.

1940. Pinon and juniper: a preliminary study of volume, growth, and yield. U.S. Soil Conserv. Serv. Region Bul. 71, Region 8, Forest Serv. 12, 86 pp.

1941. Pinon and juniper woodlands of the Southwest. Jour. Forestry 39: 542-545.

Hull, A. C., Jr., and C. W. Doran.

1950. Preliminary guide to reseeding pinyon-juniper lands of western Colorado. Rocky Mountain Forest and Range Expt. Sta. Paper No. 4

Humphrey, R. R.

1950. Arizona range resources, ii. Yavapai County, Ariz. Agr. Expt. Sta. Bul. 229, 55 pp., illus.

1953. Forage production on Arizona ranges, iii. Mohave County. Ariz. Agr. Expt. Sta. Bul. 244, 79 pp., illus.

1955. Forage production on Arizona ranges, iv. Coconino, Navajo, Apache Counties. A study in range condition. Ariz. Agr. Expt. Sta. Bul. 266, 83 pp., illus.

1958. The desert grassland - a history of vegetational change and analysis of causes. Botan. Rev. 24: 193-252. (Reprinted as Ariz. Agr. Expt. Sta. Bul. 299.)

1959. Your range--its management. Ariz. Agr. Expt. Sta. Special Rpt. 2, 35 pp., illus.

1959. Forage and water. Jour. Range Manage. 12: 164-170.

1960. Forage production on Arizona ranges, v. Pima, Pinal, and Santa Cruz Counties. Ariz. Agr. Expt. Bul. 302, 137 pp., illus.

, A. L. Brown, and A. C. Everson

1952. Common Arizona range grasses. Ariz. Agr. Expt. Sta. Bul. 243, 102 pp., illus. (Revised as Ariz. Agr. Expt. Sta. Bul. 298.)

Humphrey, R. R., and R. J. Shaw

1957. Paved drainage basins as a source of water for live-stock and game. Jour. Range Managt. 10: 59-62.

Huss, Donald L.

1954. Factors influencing plant succession following fire in ash-juniper types in Real County, Texas. Thesis, A & M College of Tex., 80 pp.

Illige, Denis J.

1954. Ecological investigation of Study Area No. 3. Ariz. Game and Fish Comm. Proj. W-71-R-1, Work Plan 4, Job 3. ✓

Jackson, C. W.

1928. Seed germination in certain New Mexico range grasses. Botan. Gaz. 86: 270-294.

Jaenicke, Alexander J., and Max H. Foerster

1915. The influence of western yellow pine forest on the accumulation and melting of snow. U.S. Monthly Weather Rev. 43: 115-126.

Jameson, Donald A.

1957. Range management research at Flagstaff Research Center, Rocky Mountain Forest and Range Expt. Sta. (a project analysis and working plan). 65 pp.

Jeffers, D. S.

1921. The pinon-juniper land problem I: should the pinon-juniper lands be included in the National Forests. Jour. Forestry 19: 534-537.

Jenkins, R. B.

1939. Cedar and poverty. The Cattleman 28(1): 61-64.

Johnsen, T. N., Jr.

1959. Junipers (Handbook: chemical control of range weeds). pp. W9-W-10.

-
1959. Longevity of stored juniper seeds. Ecology 40: 487-488.

Johnsen, Thomas N., Jr.

1962. One-seed juniper invasion of northern Arizona grasslands. Ecol. Monog. 32: 187-209.

Johnson, W. M.

1953. Effect of grazing intensity upon vegetation and cattle gains on ponderosa pine-bunchgrass ranges of the Front Range of Colorado. U.S. Dept. Agr. Cir. 929, 36 pp.

Johnson, W. M.

1956. The effect of grazing intensity on plant composition, vigor, and growth of pine-bunchgrass ranges in central Colorado. Ecology 37(4): 790-798.

- _____, 1959. Grazing intensity trials on seeded ranges in the ponderosa pine zone of Colorado. Jour. Range Mangt. 12: 1-7.

_____, and Elbert H. Reid

1958. Herbage utilization on pine-bunchgrass ranges of Colorado. Jour. Forestry 56(9): 647-651.

_____, J. G. Cady, and M. S. James

1959. Characteristics of some brown grumusols of Arizona. Paper presented to the Amer. Soc. Agron.

_____, and C. H. Niederhof

1941. Some relationships of plant cover to runoff, erosion, and infiltration on granitic soils. Jour. Forestry 39: 854-858.

Kaeiser, Margaret

1953. Macrostructure of the wood of junipers. Botan. Gaz. 115: 155-162.

Karmysheva, N. Kh.

1962. Spread of juniper in the national forest. Iz Vest. Akad Nauk, Kazakhsk. SSR. Ser Bot i Pochvovedenie 1. 13: 96-104.

Kinbacher, E. J., and M. Laude Horton

1955. Frost heaving of seedlings in the laboratory. Agron. Jour. 47: 415-418.

Kincaid, David Reed

1959. The upper desert grassland of southern Arizona--a basic ecological analysis. MS Thesis, Univ. of Ariz.

Kincaid, D. R., and Others.

1959. The spread of Lehmann lovegrass as affected by mesquite and native perennial grasses. Ecology 40: 738-742.

Kneebone, William R.

1957. Blue grama seed production studies. Jour. Range Mangt. 10: 17-21.

- Kneebone, William R., and Carlos L. Cremer
1955. The relationship of seed size to seedling vigor in some native grass species. Agron. Jour. 47: 472-477.
- _____, and R. W. Greve
1961. Economic potentials from blue grama (Bouteloua gracilis) seed production under irrigation in northwest Oklahoma. Jour. Range Mangt. 14(3): 138-143.
- Kozlowski, Theodore T.
1964. Shoot growth in woody plants. Botan. Rev. 30: 335-392.
- Krauch, Hermann
1949. Results of thinning experiment in ponderosa pine pole stands in central Arizona. Jour. Forestry 47: 466-469.
- Krefting, L. W.
1941. Methods of increasing deer browse. Jour. Wildl. Mangt. 5: 95-102.
- Kucera, C. L., John H. Ehrenreich, and Carl Brown.
Some effects of fire on tree species in Missouri prairie. Iowa State Jour. Sci. 38: 179-185.
- Lantow, J. R., and others
1943. Range livestock production practices in the Southwest. U.S. Dept. Agr. Misc. Publ. 529, 21 pp., illus.
- Laude, Horton M.
1958. Use of herbicides in competition studies of range vegetation. Ecology 39: 537-538.
- Lavin, Fred
1943. Variations in the responses of different geographic strains of blue grama grass to photoperiod. Unpub. Ph.D. dissertation, Univ. of Chicago.
- _____, and H. W. Springfield
1955. Seeding in the southwestern pine zone for forage improvement and soil protection. U.S. Dept. Agr. Agr. Handbook 89, 52 pp., illus.
- Leiberg, J. G., T. F. Rixon, and A. Dodwell
1904. Forest conditions in the San Francisco Mountain Forest Reserve, Arizona. U.S.G.S. Professional Paper 22, 95 pp.

Leigh, J. H.

1961. The unity of grass stool in Eragrostis curvula. Jour. So. Afr. Botan. 27: 99-100.

Lemon, E. R.

1962. Soil aeration and plant root relations. I. Theory. Agron. Jour. 54: 167-170.

_____, and C. C. Wiegand

1962. Soil aeration and plant root relations. II. Root respiration. Agron. Jour. 54: 171-175.

Lemon, P. C.

1946. Prescribed burning in relation to grazing in the longleaf slash pine type. Jour. Forestry 44: 115-117.

Leopold, A.

1924. Grass, brush, timber and fire in southern Arizona. Jour. Forestry 22: 1-10.

Leopold, L. B.

1951. Vegetation of southwestern watersheds in the nineteenth century. Geog. Rev. 41: 295-316.

Lillie, D. T.

1962. Herbicide combinations for the control of shrub live oak. Res. Prog. Rpt., West. Weed Control Conf. Proc. 1962: 17-18.

- _____. 1963. Control of Arizona chaparral with 2,4,5-T and silvex. Jour. Range Mangt. 16: 195-199, illus.

_____, George E. Glendening, and C. P. Pase

1964. Sprout growth of shrub live oak as influenced by season of burning and chemical treatments. Jour. Range Mangt. 17: 69-72, illus.

Lindenmuth, A. W., Jr.

1960. A survey of effects of intentional burning on fuels and timber stands of ponderosa pine in Arizona. U.S. Forest Serv. Rocky Mountain Forest and Range Expt. Sta. Sta. Paper 54, 22 pp.

_____, and James R. Davis

1962. Chemical treatment helps burn brush safely in Arizona. The Timberman 43: 9.

Lindenmuth, A. W., Jr., and G. E. Glendening

1962. Controlled burning of Arizona chaparral a 1962 progress report. Sixth Ann. Watershed Symp. Proc., Sept. 18, 1962. pp. 23-24.

Lindsey, A. A.

1949. Factors influencing the extension of conifers to unusually low elevation on a New Mexico lava bed. Bul. Ecol. Soc. Amer. 39(4): 58-59.

Linnartz, N. E.

1954. Burning cedar harmful. Soil and Water Mag. 3(12): 24-25.

Lister, P. B.

1938. How utilization of forage plants varies by seasons on semidesert grasslands. Southwestern Forest and Range Expt. Sta. Res. Note 40, 3 pp.

1938. Seasonal utilization of forage plants. Cattleman, 29-30, Aug.

_____, and F. X. Schumacher

1937. The influence of rainfall upon tuft area and height growth of three semi-desert range grasses in southern Arizona. Jour. Agr. Res. 54: 109-121.

Little, Elbert L, Jr.

1937. Harvesting the Southwest's pinon nut crop. (Flagstaff file copy)

1938. Alligator juniper. Southwestern Forest and Range Expt. Sta. Res. Note 30, 2 pp.

1938. Stages of growth on pinons in 1938. Southwestern Forest and Range Expt. Sta. Res. Note 50, 4 pp. Dec.

1938. The earliest stages of pinon cones. Southwestern Forest and Range Expt. Sta. Res. Note 46, 4 pp., Dec.

1941. Managing woodlands for pinon nuts. Chron. Botan. Vol. VI, 15, pp. 348-349.

- Little, Elbert L., Jr.
1943. Common insects on pinyon (Pinus edulis). Reprint from Jour. New York Ent. Soc. Vol. LI, pp. 239-252.
-
1950. Southwestern trees. U.S. Dept. Agr. Agr. Handbook No. 9, 109 pp., illus.
- Livingston, R. B.
1947. An ecological survey of the Black Forest region and adjacent plains. Ph.D. Thesis, Duke Univ., Durham, N. C.
- Loehwing, W. F.
1937. Root interactions of plants. Botan. Rev. 3: 195-239.
- Loftfield, J. V. G.
1923-24. Grazing ranges of northern Arizona. Carnegie Inst. of Wash. Yearbook 23: 264-265.
- Longhurst, W. M.
1956. Stump sprouting of oaks in response to seasonal cutting. Jour. Range Mangt. 9(4): 194-196.
- Lowe, Charles H., Jr.
1961. Biotic communities in the sub-Mogollon region of the inland Southwest. Jour. Ariz. Acad. Sci. 2: 40-49.
- Lucas, H. L., and G. O. Mott
1962. Methods of computing results of grazing trials. Range Res. Methods Conf., Denver, May 7-12. Vol. I, Sec. II. I.
- Lunt, Herbert A.
1951. Liming and twenty years of litter raking and burning under red (and white) pine. Soil Sci. Soc. Proc. 15.
- Lupe, Nelson
1958. Juniper eradication on the Fort Apache Indian Reservation. Abstracts of papers presented at the 11th Ann. meeting, Amer. Soc. of Range Mangt., p. 17.
- Magee, A. C.
1957. Goats pay for clearing Grand Prairie rangelands. Tex. Agr. Expt. Sta. MP-206, 8 pp.

- Mann, Dudley T., and Robert S. Hays
1948. Effect of grass on invasion of cedar. Jour. Soil and Water Conserv. 3(1): 49.
- Martin, S. Clark, and John S. Crosby
1955. Burning and grazing on glade ranges in Missouri Central States Forest Expt. Sta. Tech. Paper 148, 13 pp.
- Martin, W. P.
1940. Distribution and activity of azotobacter in the range and cultivated soils of Arizona. Ariz. Agr. Expt. Sta. Tech. Bul. 83: 335-369.
- _____, and J. E. Fletcher
1943. Vertical zonation of great soil groups on Mt. Graham, Arizona, as correlated with climate, vegetation, and profile characteristics. Ariz. Agr. Expt. Sta. Tech. Bul 99, 153 pp., illus.
- Martinez, Maximino
1946. Los Juniperus Mexicanos. Tom 17, Anales del Instituto de Biologia della Universidad Nacional de Mexico, D. F., 128 pp.
- McAllister, Dean F.
1943. The effect of maturity on the viability and longevity of the seeds of western range pasture grasses. Jour. Amer. Soc. Agron. 35: 442-453.
- McCulloch, Clay
1954. Ecological investigation of Three Bar Study Area. Ariz. Game and Fish Comm. Proj. W-71-R-1, Work Plan 4, Job 1, 31 pp.
- McCulloch, C. Y.
1955. Field observations of deer in the Three-Bar vicinity. Ariz. Game and Fish Dept., Comp. Rpt., Proj. W-71-R2, 24 pp.
- McCulloch, Clay Y., Jr.
1955. Utilization of winter browse on wilderness big game range. Jour. Wildl. Mangt. 19: 206-215.
- McCulloch, Clay Y.
1962. Watershed and game management. 6th Ann. Watershed Symp. Proc., Ariz., pp. 25-27.

McDonald, James E.

1956. Variability of precipitation in an arid region: a survey of characteristics for Arizona. Univ. of Ariz. Inst. of Atmos. Physics, Tech. reports of the Met. and climatology of arid regions No. 1, 88 pp.

McDougall, W. B., and Margaret C. Jacobs

1927. Tree mycorrhizas from the central Rocky Mountain region. Amer. Jour. Botan. 14: 258-266.

McGinnies, W. G., and J. F. Arnold

1939. Relative water requirement of Arizona range plants. Ariz. Agr. Expt. Sta. Tech. Bul. 80: 167-246.

_____, K. W. Parker, and G. E. Glendening

1941. Southwestern range ecology. Southwestern Forest and Range Expt. Sta., 211 pp., illus.

McKee, E. D.

1936. The environment and history of the Toroweap and Kaibab-formations of northern Arizona and southern Utah. Carnegie Inst. of Wash. Publ. 492.

McLaren, A. D.

1963. Biochemistry and soil science. Science 141: 1141-1147.

McMahan, C. A.

1964. Comparative food habits of deer and three classes of livestock. Jour. Wildl. Mangt. 28: 798-808.

Mead, P.

1930. An ecological description of the Kaibab Plateau, Arizona. Unpubl. Master's Thesis, Univ. of Chicago.

Meagher, G. S.

1943. Reaction of pinyon and juniper seedlings to artificial shade and supplemental watering. Jour. Forestry 41: 480-482.

- _____. 1948. The 1948 pinyon nut crop in Arizona and New Mexico. Southwestern Forest and Range Expt. Sta. Note 116.

Meagher, George

1950. Ponderosa pine research in the Southwest (A problem analysis.) Typed mss., 60 pp.

- Manzies, J. D.
1964. Survival of microbial plant pathogens in soil.
Botan. Rev. 29: 79-122.
- Merkle, J.
1952. An analysis of a pinyon-juniper community at Grand Canyon, Arizona. Ecology 33: 375-384.
- Merriam, C. H.
1890. Results of biological survey of the San Francisco Mountain Region and the desert of the Little Colorado, Arizona. No. Amer. Fauna 3.
- Merrick, G. D.
1937. The effect of grazing on the density of forage grasses on the Coconino Plateau in northern Arizona. Thesis, Duke Univ., 33 pp. (Copy on file, Rocky Mountain Forest and Range Expt. Sta., Flagstaff, Ariz.)
- Miller, Fred H.
1921. Reclamation of grasslands by Utah juniper on the Tusayan National Forest, Arizona. Jour. Forestry 19: 64.
- Moore, Thomas C.
1963. A germination inhibitor in the achenes of Cercocarpus montanus. Ecology 44: 406-409.
- Muller, C. H.
1953. The association of desert annuals with shrubs. Amer. Jour. Botan. 40: 53-60.
- _____, W. H. Muller, and B. L. Haines.
1964. Volatile growth inhibitors produced by aromatic shrubs. Science 143: 471-473.
- Muller, W. H., and C. H. Muller
1956. Association patterns involving desert plants that contain toxic products. Amer. Jour. Botan. 43: 354-361.
- Murie, Adolph.
1951. Coyote food habits on a southwestern cattle range. Jour. Mammalogy 32: 291-295.
- Myers, Clifford A.
1962. Twenty year growth of Utah juniper in Arizona. U.S. Forest Serv. Rocky Mountain Forest and Range Expt. Sta. Res. Note 71, 2 pp.

- Nakayama, F. S., and A. D. Scott
1961. Gas sorption by soils and clay minerals: 1. Solubility of oxygen in moist materials. Soil Sci. 94(2): 106-110.
- Neel, R. B., and J. S. Olson.
1962. The use of analog computers for simulating the movement of isotopes in ecological systems. Oak Ridge Natl. Lab. ORNL-3172.
- Neff, D. J.
1960. Deer population trend techniques. Proj. w-78-u-4. Work Plan 1, Job 4. Completion report May.
- Nichol, A. A.
1952. The natural vegetation of Arizona. Ariz. Univ. Agr. Expt. Sta. Bul. 127: 187-230, illus.
- Odum, H. T.
1960. Ecological potential and analog circuits for the ecosystem. Amer. Sci. 48: 1-8.
- Olmstead, C. E.
1952. Photoperiodism in native range grasses. 6th Internatl. Grassland Cong. Proc. 1952: 676-682.
- Olsen, C.
1961. Competition between trees and herbs for nutrient elements in calcareous soil. Soc. Expt. Biol. Symp. 15: 145-155.
- Olsen, C. F., and E. J. Dyksterhuis
1940. A study of cedar in west central Texas with special reference to its watershed protection value. No. Concho Survey Spec. Invest. Rpt. 34 pp.
- Olson, J. S., and D. A. Crossley, Jr.
1963. Trace studies of the breakdown of forest litter. In Radioecology, Vincent Schultz and Alfred W. Klement, Jr., Eds. Reinhold Publ. Co., and the Amer. Inst. of Biol. Sci.
- Oosting, H. J., and P. J. Kramer
1946. Water and light in relation to pine reproduction. Ecology 27: 47-53.
- Palmer, E. J.
1920. The canyon flora of the Edwards Plateau of Texas. Jour. Arnold Arboretum 1: 233-239.

Parker, Kenneth W.

1939. The control of snakeweed in the Southwest. Southwestern Forest and Range Expt. Sta. Res. Note 76, 4 pp.

Parker, K. W.

1939. Prevention of death losses from pinque. Natl. Wool Grower, 18-19. Nov.

1939. Management of livestock to avoid losses from poisonous plants. Southwestern Forest and Range Expt. Sta. Res. Note 70, 4 pp.

1939. Control of noxious plants in the Southwest. Southwestern Forest and Range Expt. Sta. Res. Note 77, 4 pp.

Parker, Kenneth W.

1942. General guide to satisfactory utilization of the principal southwestern range grasses. Southwestern Forest and Range Expt. Sta. Res. Note 104, 4 pp.

Parker, K. W.

1943. Market 'em when they're ready. Cattleman, 2 pp. Sept.

1945. Juniper comes to the grasslands. Amer. Cattle Prod., 4 pp. Nov.

1945. Forest grazing in the Southwest. Soc. Amer. Foresters Proc., 278-286.

1948. An analysis of range problems in the Southwest, 125 pp.

1952. New trends in standards of range use. Jour. Forestry 50: 856-859.

, and G. E. Glendening

1942. General guide to satisfactory utilization of the principal southwestern range grasses. Southwestern Forest and Range Expt. Sta. Res. Note 104, 4 pp.

Pase, G. P.

1958. Herbage production and composition under immature ponderosa pine stands in the Black Hills. Jour. Range Mangt. 11: 238-243.

Pase, Charles P., and Richard M. Hurd

1957. Understory vegetation as related to basal area, crown cover and litter produced by immature ponderosa pine stands in the Black Hills. Soc. Amer. Foresters Proc., pp. 156-158.

_____, and Floyd W. Pond

1964. Vegetation changes following the Mingus Mountain burn. U.S. Forest Serv. Res. Note RM-18, 8 pp.

Paulsen, Harold A., Jr., and Fred N. Ares

1961. Trends in carrying capacity and vegetation on an arid southwestern range. Jour. Range Mangt. 14: 78-83.

_____, and Fred N. Ares

1962. Grazing values and management of black grama and tobosa grasslands and associated shrub ranges of the Southwest. U.S. Dept. Agr., Forest Serv. Tech. Bul. 1270, 56 pp.

Pearson, G. A.

1931. Forest types in the Southwest as determined by climate and soil. U.S. Dept. Agr., Tech. Bul. 247, 144 pp.

_____. 1936. Why the prairies are treeless. Jour. Forestry 34(4): 405-408.

_____. 1940. Forest land use. Jour. Forestry 38: 261-270.

_____. 1944. Multiple use in forestry. Jour. Forestry 42: 243-249.

_____. 1948. Retired forester's views on timber vs. grass issue. Ariz. Farmer, Jan. 10, pp. 24-25.

_____. 1950. Management of ponderosa pine in the Southwest. U.S. Dept. Agr. Agr. Monog. No. 6: 218 pp.

Pessin, L. J.

1938. The effects of vegetation on the growth of longleaf pine seedlings. Ecol. Monog. 8: 115-149.

_____, and R. A. Chapman

1944. The effect of living grass on the growth of longleaf pine seedlings in pots. Ecology 25: 85-90.

Phillips, F. J.

1909. A study of pinon pine. Bot. Gaz. 48: 216-223.

1910. The dissemination of juniper seeds by birds. Forestry Quart. 8: 60-73.

_____, and W. Mulford

1912. Utah juniper in central Arizona. U.S. Dept. Agr. Forest Serv. Cir. 197, 19 pp.

Pickett, G. S.

1955. Report of burning of Grayer Wash juniper burning project, Walnut Creek District, Prescott National Forest. Unpubl. rpt., 7 pp.

Pickrell, K. P., and E. B. Stanley

1930. An economic study of range sheep production in Arizona. Ariz. Agr. Expt. Sta. Bul. 134: 519-551.

Pingrey, H. B., and E. J. Dortignac

1959. Economic evaluation of seeding crested wheatgrass on northern New Mexico rangeland. N. Mex. Agr. Expt. Sta. Bul. 433, 80 pp., illus.

Pistor, W. J.

1952. Larkspur. Ariz. Cattlelog, 64-66. June.

1952. Poisoning by pigweed or careless weed. Ariz. Cattlelog, 55-56. July.

1958. Nitrate (nitrite) poisoning. Ariz. Cattlelog, p. 63, Dec.

Plummer, A. P.

1943. The germination and early seedling development of 12 range grasses. Jour. Amer. Soc. Agron. 35: 19-33.

Plummer, A. Perry

1958. Restoration of juniper pinyon ranges in Utah. Soc. Amer. Foresters Proc. 1958: 207-211.

Plummer, F. G., Theodore P. Rixon, and Arthur Dodwell

1904. Forest conditions in the Black Mesa Forest Reserve, Arizona. U.S. Geol. Survey Prof. Paper 23.

Potter, Loren D.

1957. Phytosociological study of San Augustin Plains, New Mexico. Ecol. Monog. 27: 115-136.

Potter, Loren D., and John C. Krenetsky

Plant succession with released grazing on New Mexico range lands.

Randles, Quincy

1949. Pinyon-juniper in the Southwest. U.S. Dept. Agr. Yearbook 1949: 342-347.

Rasmussen, D. Irvin

1941. Biotic communities of Kaibab Plateau, Arizona. Ecol. Monog. 11: 229-275.

Rauzi, F., and R. L. Lang

1956. Improving shortgrass range by pitting. Wyo. Agr. Expt. Sta. Bul. 344, 11 pp.

Read, A. D.

1915. The flora of the Williams Division of the Tusayan National Forest, Arizona. Plant World 18: 112.

Reid, Elbert H.

1964. Forage production in ponderosa pine forests. Soc. Amer. Foresters Proc. 1964: 61-64.

Reynolds, H. G.

1959. Managing grass-shrub cattle ranges in the Southwest. U.S. Dept. Agr. Agr. Handbook 162, 40 pp., illus.

1960. Watershed management research in Arizona and New Mexico. Jour. Forestry 58: 275-278.

Reynolds, Hudson G.

1962. Effect of logging on understory vegetation and deer use in a ponderosa pine forest of Arizona. U.S. Forest Serv. Rocky Mountain Forest and Range Expt. Sta. Res. Notes 80, 7 pp.

1962. Selected bibliography on range research in Arizona. Ariz. Sect. Soc. Range Mangt. Proc., pp. 50-68.

1962. Some characteristics and uses of Arizona's major plant communities. Jour. Ariz. Acad. Sci. 2: 62-71.

- Reynolds, H. G., F. Lavin, and H. W. Springfield
1949. Preliminary guide for range reseeding in Arizona and New Mexico. Southwestern Forest and Range Expt. Sta. Res. Rpt. 7, 14 pp., illus.
- _____, and G. E. Glendening
1959. Research in management of chaparral lands in Arizona. Ariz. Cattlelog 14(12): 13-16.
- _____, and H. W. Springfield
1953. Reseeding southwestern range lands with crested wheatgrass. U.S. Dept. Agr. Farmers' Bul. 2056, 20 pp.
- Reveal, J. L.
1944. Single leaf pinyon and Utah juniper woodlands of western Nevada. Jour. Forestry 42: 276-278.
- Rice, Elroy L.
1950. Growth and floral development of five species of range grasses in central Oklahoma. Botan. Gaz. 111: 361-377.
- Ribb, L. R.
1952. Consumptive use of water--forest and range vegetation. Trans. Amer. Soc. Civ. Engin. 117: 974-990
- _____
1956. Relation of plant cover to watershed management. Ariz. Cattlelog 11(12): 34-36.
- _____
1962. Erosion and sediment movement following a wildfire in a ponderosa pine forest of central Arizona. U.S. Forest Serv. Rocky Mountain Forest and Range Expt. Sta. Res. Notes 76, 12 pp.
- Riedl, W. A., R. L. Lang, and K. Asay
1959. Selection and improvement of Eurotia lanata (winterfat) and other species for salt-desert areas. Wyo. Agr. Expt. Sta. Mimeo. C. 114: 22-28. June.
- Riewe, Marvin E.
1961. Use of the relationship of stocking rate to gain of cattle in an experimental design for grazing trials. Agron. Jour. 53: 309-313.
- Rigden, J. T.
1939. Facts concerning Arizona ranching. Ariz. Agr. Exp. Serv., 7 pp.

- Rigden, J. T., and K. W. Parker
1943. Guides to condition of Arizona chaparral goat ranges.
Ariz. Stockman. 4, 10, 11, 13, 15. Nov.
- Roberts, N. K.
1963. Economic foundations for grazing use fees on public
lands. Jour. Farm. Econ. 45: 721-731.
- Robinson, H. H.
1913. The San Franciscan volcanic field, Arizona. U.S.
Geol. Survey Prof. Paper 76, 213 pp.
- Rosenberg, Norman J., and N. A. Willits
1962. Yield and physiological response of barley and beans
grown in artificially compacted soils. Soil Sci.
Soc. Amer. Proc., Vol. 26, No. 1. Jan-Feb.
- Rovira, A. D.
1962. Plant-root exudates in relation to the rhizosphere
microflora. Soils and Ferts 25, No. 3, 167-172.
- Sampson, Arthur W.
1944. Plant succession on burned chaparral lands in northern
California. Univ. Calif. Col. Agr. Bul. 685.
- Sampson, A. W.
1951. A symposium on rotation grazing in North America.
Jour. Range Mangt. 4: 19-24.
- _____, and Beryl S. Jespersen
1964. California brushlands and range browse plants. Univ.
of Calif. Div. Agr. Brushland Manual, 162 pp.
- Schaus, Richard
1951. It's a battle--Arizona's native grass is fighting
juniper to live. West. Livestock Jour. 29:1100, 101,
103. Oct.
- Schaus, R. G.
1952. Juniper eradication in the west. Amer. Cattle Prod.
pp. 11, 12. Nov.
- Schmutz, E. M., D. R. Cable, and J. J. Warwick
1959. Effects of shrub removal on vegetation of semi-desert
grass-shrub ranges. Jour. Range Mangt. 12: 34-37.
- Schmutz, Erwin M., and Raymond M. Turner
1957. Herbicide tests on fire sprouts of turbinella oak in
the Arizona chaparral. Res. Prog. Rpt., West. Weed
Control Conf. Proc. 1957: 44.

- Schmutz, E. M., and D. W. Whitman
1962. Shrub control studies in the oak-chaparral of Arizona,
Jour. Range Mangt. 15: 61-67.
- Schoffelmayer, F. H.
1939. Texas problem, grass vs. cedars. Cattleman, p. 38.
- Schram, J. R.
1958. The mechanism of frost heaving of tree seedlings.
Amer. Phil. Soc. Proc. 102: 333-351.
- Schroeder, Bill
1958. Controlled burning on the Hualapai Reservation. Ariz...
Cattlelog, p. 50. May.
- Schubert, Gilbert H.
1963. Natural regeneration in the Southwest--how do we get
it? how much can we depend on it? Files,
Rocky Mountain Forest and Range Expt. Sta.,
Flagstaff, Ariz.
- _____
1964. Forest management research of ponderosa pine in the
Southwest. Files, Rocky Mountain Forest and Range
Expt. Sta., Flagstaff, Ariz.
- _____, L. J. Heidmann, and M. M. Larson
1964. Artificial reforestation practices for the Southwest.
(Submitted for U.S. Dept. Agr. publication.)
- Schultz, A. M., J. L. Launchbaugh, and H. H. Biswell
1955. Relationship between grass density and brush seedling
survival. Ecology 36: 226-238.
- Schultz, E. F., Jr., W. R. Langford, E. M. Evans, and others
1959. Relationship of beef gains to forage yields. Agron.
Jour. 51: 207-211.
- Schuster, J. L., and L. K. Halls
1962. Timber overstory determines deer forage in shortleaf-
loblolly pine-hardwood forests. Soc. Amer. Foresters
Proc., pp. 165-167.
- Scott, T. W.
1961. The influence of several physical soil factors on the
development and activity of plant roots. Disser.
Abstr. 21(12): 3586. June.
- Searles, R. D.
1951. The drought in southwestern United States. U.S. Dept.
Int., 65 pp., illus.

- Sellers, William D.
1960. Arizona climate. Univ. of Ariz. Press, Tucson.
- Sen Gupta, M. B., and A. H. Cornfield
1963. Phosphorus in calcareous soils. III. "Available" phosphate in calcareous soils as measured by five chemical methods and phosphate uptake by fye-grass in a pot test. Jour. Sci. Food Agr. 14: 563-567.
- Shannon, Stanton
1953. The effects of forest fires or of controlled burning on the forest soils of northern Arizona. A thesis, 58 pp.
- Shantz, H. L., and Rafael Zon
1924. The natural vegetation of the United States. U.S. Dept. Agr., Bur. Agr. Econ., Atlas Amer. Agr. pt. 1, sec. E., 29 pp., illus.
- Shirley, Hardy L.
1945. Light as an ecological factor and its measurement. II. Botan. Rev. 11: 497-532.
- Shreve, F.
1942. The vegetation of Arizona. In Flowering plants and ferns of Arizona. U.S. Dept. Agr. Misc. Publ. 423: 10-23, illus.
- Skau, Clarence McClelland
1960. Some hydrologic characteristics in the Utah juniper type of northern Arizona. Thesis for the degree of Ph.D., Michigan State Univ.
- Smith, Arthur D.
1950. Inquiries into differential consumption of juniper by mule deer. Utah Fish and Game Bul. (Oct.): 4.
- Smith, A. D.
1952. Digestibility of some native forage for mule deer. Jour. Wildl. Mangt. 10: 309-312.
- Smith, Arthur D., and Phillip J. Urness.
1962. Analyses of the twig length method of determining utilization of browse. Utah State Dept. Fish and Game Publ. 62-9, 35 pp.
- Smith, H. V.
1956. The climate of Arizona. Ariz. Agr. Expt. Sta. Bul. 279, 99 pp.

Smith, J. G.

1899. Grazing problems in the Southwest and how to meet them.
U.S. Dept. Agr. Bul. 16 (Agros. 32), 46 pp.

Smoliak, S.

1956. Influence of climatic conditions on forage production
of shortgrass rangeland. Jour. Range Mangt. 9: 89-91,
illus.

_____, and H. F. Peters

1955. Climatic effects on foraging performance of beef cows
on winter range. Canad. Jour. Agr. Sci. 35: 213-216.

Sneva, Forrest A., and D. N. Hyder

1962. Estimating herbage production on semiarid ranges in the
Intermountain region. Jour. Range Mangt. 15: 88-93,
illus.

Society of American Foresters

1954. Forest cover types of North America (exclusive of
Mexico). Wash., D. C., Soc. Amer. Foresters, 67 pp.

Stagner, Marilyn L.

1962. Standing crop and primary productivity in the tree
stratum of a pinyon-juniper community in northern
Arizona. Bul. Ecol. Soc. Amer. 43(3): 90-91.

Stanley, E. B.

1938. Nutritional studies with cattle on a grassland-type
range in Arizona. Ariz. Agr. Expt. Sta. Tech. Bul.
79: 129-164.

_____, and C. W. Hodgson

1938. Seasonal changes in the chemical composition of some
Arizona range grasses. Ariz. Agr. Expt. Sta. Tech.
Bul. 73: 451-466.

Stoddart, L. A., and J. E. Greaves

1942. The composition of summer range plants in Utah. Utah
Agr. Expt. Sta. Bul. 305, 22 pp.

Stotsky, G.

1962. Some effects of clay minerals on microbial ecology in
soil. VIIIth Internatl. Cong. Microbiology, Biol.
Abstr. 41: 3537.

Stubblefield, T. M.

1954. Cow-calf versus cow-yearling production program for
the Southwest. Ariz. Agr. Expt. Sta., 11 pp.

Stubblefield, T. M.

1956. Greater returns from cow-yearling operations on Southwest ranges. Jour. Range Managt. 9: 8-10.

Sudworth, G. B.

1915. The cypress and juniper trees of the Rocky Mountain region. U.S. Dept. Agr. Bul. 207, 36 pp.

Sumner, C. F.

1960. Germination inhibitors in plant material. In Plant Phenolics Group. Phenolics in plants in health and disease, pp. 25-34. Pergamon Press, New York.

Summer, D. C., and Rodney D. Cobb

1962. Postharvest dormancy of Coronado side-oats grama Bouteloua curtipendula (Michx.) Torr. as affected by storage temperature and germination inhibitors. Crop. Sci. 2: 321-325.

Swank, W. G.

1956. Protein and phosphorus content of browse plants as an influence on southwestern deer herd levels. Trans. 21st No. Amer. Wildl. Conf., pp. 141-158.

1958. The mule deer in Arizona chaparral. Ariz. Game and Fish Dept., Wildl. Bul. 3, 109 pp.

Talbot, M. W.

1937. Indicators of southwestern range conditions. U.S. Dept. Agr. Farmers' Bul. 1782, 36 pp., illus.

Tarrant, Robert F.

1957. Soil moisture conditions after chemically killing manzanita brush in central Oregon. U.S. Forest Serv. Pacific Northwest Forest and Range Expt. Sta. Res. Note 156, 4 pp.

Taylor, W. P., and J. V. G. Loftfield

- 1923-24. Rodent damages on Arizona ranges. Carnegie Inst. of Wash. Yearbook 23: 265.

_____, C. T. Vorhies, and P. B. Lister

1935. The relation of jackrabbits to grazing in southern Arizona. Jour. Forestry 33: 490-498.

Thomsen, F. L.

1954. More efficient marketing of Arizona cattle. Ariz. Agr. Expt. Sta. Bul. 254, 39 pp.

Thornthwaite, C. W. et al.

1942. Climate and accelerated erosion in the arid and semi-arid Southwest, with special reference to the Polacca Wash drainage basin, Arizona. U.S. Dept. Agr. Tech. Bul. 808, 134 pp., illus.

Toustenko, V. K.

1954. Scrub juniper, reserves of raw material. Lesn. Khoz. 7(2): 80-84.

Tribe, D. E.

1950. The behavior of the grazing animal: a critical review of present knowledge. Jour. Brit. Grassland Soc. 5: 209-244.

Trosko, I. K., and I. G. Veselov.

1954. Regeneration of juniper forests in the mountains of central Asia. Lesn. Khov. 5: 66-70.

Turner, Harry C.

1913. The germination of alligator juniper seed. Rev. Forest Serv. Invest. 2: 49-52.

Twitchell, LaForest T.

1955. Germination of fourwing saltbush seed as affected by soaking and chloride removal. Jour. Range Mangt. 8: 218-220.

Upton, Arthur

1937. Does extra rainfall always make more grass? Southwestern Forest and Range Expt. Sta. Res. Note 11, 3 pp.

Upton, A., W. J. Cribbs, and E. B. Stanley

1937. Occurrence of shrubs on range areas in southeastern Arizona. Ariz. Agr. Expt. Sta., 29 pp., illus.

U.S.D.A. Forest Service

1955. Timber Resource review. Chap. II., 70 pp.

-
1957. Annual report. Rocky Mountain Forest and Range Expt. Sta., pp. 89-92.

-
1958. Range plants of Arizona and New Mexico. Names, symbols, and notations, Rocky Mountain Forest and Range Expt. Sta., 86 pp.

-
1958. Timber resources for America's future. Forest Res. Rpt. 14, p. 117.

U.S.D.A. Forest Service

1963. Silvics of forest trees of the United States. Agr. Handbook 271: 398-403.
- Valentine, Kenneth A.
1938. Two proper-use factors for tobosa grass. Southwestern Forest and Range Expt. Sta. Res. Note 38, 4 pp.
- Van Haverbeke, David F.
1963. Root development of ponderosa pine seedlings in the Black Hills. Ecology 44(1): 161-165.
- Van Melle, P. J.
1953. Monoecism in Juniperus scopulorum. Phytologia 4: 172.
- Veatch, J. O.
1920. Description of the soils of sample plots in the Coconino and Tusayan National Forests, Arizona, also, the soils of the San Francisco Mountain region, Arizona. Unpub. Rpt., U.S. Dept. Agr. Bur. Soils.
- Viro, P. J.
1963. Factorial experiments in forest humus decomposition. Soil.Sci. 95: 24-30.
- Vishen, Stephen Sargent
1954. Climatic atlas of the United States. Harvard Univ. Press.
- Vorhies, C. T., and W. P. Taylor
1933. The life histories and ecology of jackrabbits, Lepus alleni and Lepus californicus, esp., in relation to grazing in Arizona. Ariz. Agr. Expt. Sta. Tech. Bul. 49: 471-587.
- _____, and W. P. Taylor
1940. Life history and ecology of the white-throated woodrat, Neotoma albigula albigula hartley, in relation to grazing in Arizona. Ariz. Agr. Expt. Sta. Tech. Bul. 86: 455-529.
- Wagar, J. A., and Clifford A. Myers
1958. Some factors affecting the results of direct-seeding ponderosa pine in the Black Hills. U.S. Forest Serv. Rocky Mountain Forest and Range Expt. Sta. Res. Notes 33, 6 pp.
- Wagle, Robert F., and Schmutz, Ervin M.
1963. The effect of fenuron on four southwestern shrubs. Weeds 11: 149-157.

- Wagoner, J. J.
1952. History of the cattle industry in southern Arizona.
Univ. Ariz. Social Sci. Bul. 20.
- Waldrip, William J.
1954. Brush control, grazing management, and revegetation of
areas infected with blueberry cedar. Tex. Agr.
Expt. Sta. Prog. Rpt. 1721, 5 pp.
- Waldrip, W. J., and D. G. Wilson
1952. What is being done about cedar. Sheep and Goat Raisers
32(8): 28-31.
- Walker, Charles F., and William A. Williams
1963. Responses of annual-type range vegetation to sulfur
fertilization. Jour. Range Mangt. 16: 64-69.
- Wallace, J. D., F. Hubbert, Jr., and R. J. Raleigh
1963. The response of yearling cattle on crested wheatgrass
pasture to energy, protein and sodium supplementation.
Jour. Range Mangt. 16: 1-5.
- Watkins, W. E.
1937. The calcium and phosphorus contents of important New
Mexico range forages. New Mex. Agr. Expt. Sta.
Tech. Bul. 246, 75 pp.
- _____
1939. Monthly variation in carotene content of two important
range grasses, Sporobolus flexuosus and Bouteloua
eriopoda. Jour. Agr. Res. 58: 695-699.
- _____
1943. Composition of range grasses and browse at varying
stages of maturity. New Mex. Agr. Expt. Sta.
Tech. Bul. 311, 43 pp.
- _____, and J. H. Knox
1945. The monthly protein and phosphorus contents of two
important range grasses of southern New Mexico.
Jour. Anim. Sci. 4: 297-305.
- Watson, J. R.
1912. Plant geography of north central New Mexico. Botan.
Gaz 54: 194-217.
- Watt, K. E. F.
1959. A mathematical model for the effect of densities of
attacked and attacking species on the number attacked.
Canad. Ent. 91: 129-144.

Weaver, J. E.

1958. Summary and interpretation of underground development in natural grassland communities. Ecol. Monog. 28: 55-78.

Whiting, Alfred F.

1942. Junipers of the Flagstaff region. Plateau 15: 23-31.

Williamson, R. E.

1964. The effect of root aeration on plant growth. Soil Sci. Soc. Amer. Proc. 28(): 86-90.

Wilson, Eldred D.

1962. A resumé of the geology of Arizona. Bul. 171, Univ. of Ariz. Press, Tucson, 140 pp.

Witkamp, Martin

1963. Microbial populations of leaf litter in relation to environmental conditions and decompositions. Ecology 44: 370-377.

Wootton, E. O.

1916. Carrying capacity of grazing ranges in southern Arizona. U.S. Dept. Agr. Bul. 367, 40 pp., illus.

Wolff, Simon E.

1946. Jackrabbits and juniper. Soil Conserv.

Wolff, S. W.

1948. An evaluation of some weedy Texas junipers. Soil Conserv. Serv. West. Gulf Region, 89 pp.

Woodbury, A. M.

1933. Biotic relationship of Zion Canyon, Utah. Ecol. Monog. 3: 147-245.

-
1947. Distribution of pigmy conifers in Utah and northeastern Arizona. Ecology 28: 113-126.

Woodhouse, S. W.

1853. Report on the natural history of the country passed over by the exploring expedition under command of Brev. Capt. L. Sitgreaves, U.S. Topographical Engineers, during the year 1851. In Report of an expedition down the Zuni and Colorado Rivers, by Captain L. Sitgreaves. Senate Executive Doc. No. 59, 32nd Cong., 2 Sess., 33-40.

Woodin, H. E.

1953. The pinyon-juniper association in the eastern foothills of the Rockies. Ph.D. Thesis, Purdue Univ.

Woodin, Howard E., and Alton A. Lindsey

1954. Juniper-pinyon east of the Continental Divide, as analyzed by the line-strip method. Ecology 35: 473-489.

Woods, F. W.

1960. Biological antagonisms due to phytotoxic root exudates. Botan. Rev. 26(4): 546-569.

Wootton, E. O.

1916. Carrying capacity of grazing ranges in southern Arizona. U.S. Dept. Agr. Bul. 367, 40 pp., illus.

Wright, Henry A.

1964. An evaluation of several factors to determine why Sitanion hystrix is more resistant to burning than Stipa comata. Ph.D. Thesis, Utah State Univ. Libr. Logan, Utah.

Wright, N., and L. J. Streetman

1958. Past performance and future potential of black grama for southwestern ranges. Jour. Range Mangt. 11: 207-214.

_____, and L. J. Streetman

1960. Grass improvement for the Southwest. Ariz. Agr. Expt. Sta. Tech. Bul. 143, 16 pp.

Young, Darrol P.

1952. Juniper-enemy of southwestern ranges. New Mex. Stockman 17: 38, 39, 58, 59. May.

Zinke, Paul J.

1962. The pattern of influence of individual forest trees on soil properties. Ecology 43(1): 130-133.

SUPPLEMENTAL LIST FOR LITERATURE CITED

- Albertson, F. W., G. W. Tomanek, and Andrew Riegel
1957. Ecology of drought cycles and grazing intensity on grasslands of central Great Plains. Ecol. Monog. 27: 27-44.
- Biswell, H. H., A. M. Schultz, and D. W. Hedrick
1953. Frost heaving of grass and brush seedlings on burned chamise brushlands in California. Jour. Range Mangt. 6: 172-180.
- Castetter, Edward F.
1956. The vegetation of New Mexico. New Mexico Quarterly, Vol. 26, No. 3. 257-288.
- Clary, W. P., and Peter F. Ffolliott
1966. Unpublished.
- _____, Peter F. Ffolliott, and Almer D. Zander
1966. Grouping sites by soil management areas and topography. U.S. Forest Serv., Rocky Mountain Forest and Range Expt. Sta. Res. Note RM-60, 4 pp.
- Corini, S.
1963. Lucerne "fatigue" and soil microflora. It. Annali. Microbiol. 13, 11-18. Herb. Abstr. 35(1): 467.
- Geiger, Rudolf
1965. The climate near the ground. Howard Univ. Press, Cambridge, Mass. 611 pp.
- Glover, Terrence, and Allen LeBaron
1966. Estimating forage production following pinyon-juniper control: a probabilistic approach. 13 pp.
- Green, Cristine R., and William D. Sellers
1964. Arizona climate. Univ. Arizona Press, Tucson. 503 pp.
- Fritts, Harold C.
1965. Tree-ring evidence for climatic changes in western North America. Monthly Weather Review, Vol. 93, No. 7, 421-443.
- Hastings, James Rodney
1964. Climatological data for Sonora and Northern Sinaloa. University of Arizona Institute of Atmospheric Physics. Technical Reports on the Meteorology and Climatology of Arid Regions. No. 15, 152 pp.

Lavin, Fred

1967. Fall fertilization of intermediate wheatgrass in the southwestern ponderosa pine zone. Jour. Range Mangt. Vol. 20(1): 16-21.

Nelson, Enoch W.

1934. The influence of precipitation and grazing upon black grama grass range. U.S. Dept. Agr. Tech. Bul. No. 409. 32 pp.

Pattison, Allan

1965. Synthesis of hourly rainfall data. Water Resources Research. Vol. 1, No. 4. 489-498.

Rich, Lowell R.

1961. Surface runoff and erosion in the lower chaparral zone - Arizona. U.S. Forest Serv., Rocky Mountain Forest and Range Expt. Sta. Paper No. 66, 35 pp.

_____, and Hudson G. Reynolds

1963. Grazing in relation to runoff and erosion on some chaparral watersheds of central Arizona. Jour. Range Mangt. Vol. 16, No. 6, 322-326.

Schubert, Walter J.

1965. Lignin biochemistry. Academic Press, New York and London. 131 pp.